

Microbiological Assessment of Gram-negative Bacterial Isolates from Ear, Eye, Nose and Throat among Patients Attending Aresho Advanced Medical Laboratory, Addis Ababa, Ethiopia

Estifanos T^{1*}
Mesele A¹
Meseret C¹

¹Department of Microbiology, Arsho Advanced Medical Laboratory, Ethiopia

Abstract

ENT samples from patients who referred to Arsho Advanced Medical laboratory were isolated for further microbiological assessment. Ninetyone (91) from both male and female patients comprising 42 ear, 21 nasal, 1 eye and 27 throat swabs respectively were screened between April -2017 to July-2017 and the samples were analyzed using culture technique, direct microscopy and identification of bacteria (Vtek 2 compact). Finding from the study identified twelve species of bacteria i.e *Pseudomonas aeruginosa*, *Proteus mirabilis*, *Kelebsiella oxytoca*, *Entrobacter cloacae*, *Citrobacter koseri*, *Citrobacter pauclus*, *Pseudomonas fulorescens*, *Klebsiella pneumoniae*, *Acinetobacter baumannii*, *E.coli*, *Leclerciaa decarboxylate* and *Proteus vulgaris*. *Proteus mirabilis* recorded the highest rates (30%) followed by *Klebsiella pneumoniae* (20%) which was isolated in ear, and nose swabs, while the least were recorded on *Citrobacter koseri*, *Citrobacter pauclus*, *Pseudomonas fulorescens*, *Proteus vulgaris*, *Acinetobacter baumannii*, *E.coli*, *Leclerciaa decarboxylate*, *Klebsiella pneumonia* (3.3%).

Key words: Bacteria, Species; Swab; Nasal.

Introduction

Ear, Eye, and Nose infection are among the most widespread and serious infections that compel an individual to seek medical attention. It represents some of the most common bacterial disease encountered affecting people of all ages. These infections are one of the leading causes of morbidity and mortality in critically ill patients [1]. The ear, nose, and throat are the frequent sites of infection, because they come in direct contact with the physical environment and are exposed to air borne microorganisms.

Disease of ear and nose affect the functioning of adults as well as children, often with significant impairment of the daily life of affected patients [1]. It has been predicted that with increase in global population, infection remain the most important causes of disease with upper respiratory infections causing hearing loss and learning disability particularly in children [2-4]. Ear infection such as chronic otitis (tinnitus), have serious consequences in developing countries such as retarded language development and progress in school among children [5]. Tinnitus which is now known to be the most common childhood infections, lead annually to the death of over 50,000 children under 5 years [6].

In addition, antimicrobial resistance profile of bacteria varies among population because of the difference in geography, local antimicrobial prescribing practices and prevalence of resistant bacterial strains in a

Article Information

Article Type: Research

Article Number: JAMBR113

Received Date: 19 October, 2018

Accepted Date: 01 November, 2018

Published Date: 07 November, 2018

***Corresponding author:** Mesele Admassie, Department of Microbiology, Arsho Advanced Medical Laboratory, Addis Ababa, Ethiopia. Tel: +251911473869, Email: [admassie90\(at\)gmail.com](mailto:admassie90(at)gmail.com)

Citation: Estifanos T, Mesele A, Meseret C (2018) Microbiological Assessment of Gram-negative Bacterial Isolates from Ear, Eye, Nose a Throat among Patients Attending Aresho Advanced Medical Laboratory, Addis Ababa, Ethiopia. J Appl Microb Res Vol: 1, Issu: 2 (27-32).

Copyright: © 2018 Estifanos T. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

given area [7]. So there should be up to date information on microbial resistance pattern at national and local levels to guide the rational use of the existing antimicrobial drugs.

The human ear, nose and throat (ENT) are closely related and inter connected parts of the body. Infections, diseases and health problems related to the ENT are therefore jointly studied and managed like the most other part of the human body. The ear, nose and throat were found to be colonized by a wide range of microorganisms some of which are more or less harmless under normal condition [8].

In Ethiopia particularly in the study area, there is no such type of recent data that shows the magnitude of the problem. Therefore, the aim of this study was to determine the gram-negative bacterial isolates and their drug susceptibility patterns from patients who gave ear and nose swab samples at Arsho Advanced medical Laboratory.

Objectives

General objectives

To determine the frequency of gram-negative bacterial outline and antibiotic susceptibility pattern isolated from ENT at Arsho advanced medical laboratory.

Specific objectives: To assess the gram-negative pathogens responsible for the ENT infection

To verify the antimicrobial resistance and sensitivity pattern of commonly isolated gram-negative bacteria from ENT patients.

Methods

Study area and period

A study was conducted from, April-2017 to July-2017 at Arsho Advanced Medical Laboratory, Addis Ababa, Ethiopia.

Samples collections

The throat swab, nasal (nose) swab, eye swab and aural (ear) swab were collected by using a sterile cotton swab, and then the swabs were transported with Amies transport media to the micro biology laboratory.

Culture technique

Throat swab were inoculated on blood agar plates, were as nasal (nose), eye and ear swab were inoculated on chocolate, blood and MacConkey agar plates. All the plates were incubated for 48 hours 5% CO₂ an aerobically with the exception of MacConkey agar plate that was inculcated in aerobic. The plates were examined for the growth of gram-negative bacteria and the pathogenic colonies were identified by VITEK 2 compact.

Bacteria identification test

Pure colonies of gram-negative colonies which were grown on macConkey Agar, chocolate Agar and Blood Agar were used for identification in VITEK 2 compact.

Suspension preparations for ID and AST card

Suspension preparation for ID card: First Transfer 3ml of 0.45% saline into a test tube. Select an isolated colony and

dissolve it. Mix well and check the density within dens check. Inoculum density for gram positive and gram-negative colonies should be between 0.5-0.63MCF. Then place the identification (ID) card into the test tube and then transfer the test tube into the Cassette.

Suspension preparation for AST card: First Transfer 3ml of 0.45% saline into a Test Tube. Transfer 145 micro liter for gram negative of the ID suspension into the saline test tube. Then place the AST card into the test tube and then transfer the test tube into the cassette.

Filling and loading the card into VITEK 2 system

Put all the Test Tubes containing Cards and suspension a Cassette. From computer work station print Cassette worksheet and record job ID and bar code for each card. When instrument status is OK, then press start fill button. Remove the Cassette from the loading station when the machine indicates.

Antibiotics used

The antibiotics used in this investigation were Ampicillin, Amoxicillin/Clavulanic Acid, piperacillin/Tazobactam, Cefalotin, Cefazolin, cefuroxime Axil, Cefoxitin, Cefpodoxime, Ceftazidime, Ceftriaxone, Cefepime, Gentamicin, Tobramycin, Ciprofloxacin, Levofloxacin, Tetracyclin, Nitrofurantoin and Trimethoprim/sulfamethoxazole for gram negative. The control strains were run simultaneously with the test organisms.

Data Analysis

Data were checked for completeness, manually, entered, and analyzed using SPSS version 20 statistical software and Excel.

Result

A total number of 42 ear swabs, 21 nasal swabs, 1 eye and 27 throat swabs were collected from the patients attending at Aresho Advanced Medical Laboratory. Out of these, ear swabs were positive in 20 (22 %), 10 (11 %) in nose swabs.

The ear swabs was positive in 9 (9.9 %) male and 11 (12.1 %) in female, nose swabs was positive 4 (4.4 %) was positive in male and 5 (5.5 %) in female (Table 1). The findings in this study show that twelve genera of gram negative bacteria which are *P. mirabilis*, *K. pneumoniae*, *E. coli*, *P. vulgaris*, *P. aeruginosa*, *P. fulorescens*, *K. oxytoca*, *E. cloacea*, *C. koseri*, *C. pauculus*, *Leclerciaa decarboxylata*, and *A. baumannii* were involved in various ear, and nose amongst patients examined at the study area. Out of all the organisms isolated 20 (66.67

Table 1: Sex Wise Distribution and Culture Result of ENT Patients diagnosed from April -2017 to July-2017 in Arsho Advanced Medical Laboratory.

Sample	Sex	Negative	Positive	Total examined
Ear	M	10	9	19
	F	12	11	23
Nose	M	3	4	7
	F	7	7	14
Throat	M	13	0	13
	F	14	0	14
Eye	M	1	0	1
	F	0	0	0

%) occurred in ear swabs followed by nasal 10 (33.3 %) while there is no growth in eye and throat.

Generally, the most prevalent organisms were *Proteus mirabilis* (30 %) while the least were *Citrobacter koseri*, *Citrobacter pauculus*, *Pseudomonas fulorescens*, *Acinetobacter baumannii* and *Leclerciaa decarboxylate* with (3.33 %). *Proteus mirabilis*, *K. pneumoniae*, and *K.oxytoca* were isolated in both ear and nose swabs, *E. coli*, *P.aeruginosa*, *P.fuloresce*, *E.cloacea*, *C.koseri*, *C.pauculus* and *A.baumannii* were isolated in ear only while *L.decarboxylata* and *P.vulgaris* were isolated from only nasal swabs (Table 2).

The age-related prevalence of the ENT infection in the study area showed that the highest prevalence of the bacterial infection (40 %) was amongst those aged 15-24 years, (26.67 %) 25-44 years, (23.33 %) 1-14 years and (10 %) 45-64 while no growth observed amongst <1 and >65 years aged as shown in (Table 3).

During the study period 18 different antimicrobial agents were used to test the antibiotic susceptibility patterns of the pathogenic gram-negative bacteria isolated from ENT patients. The general susceptibility profiles of bacterial isolates are shown in table 4. Out of the total antibiotics examined during the study period, Ampicillin had the highest overall resistance rate (80.8 %) followed by Cephalothin (58.6.6 %), Cefazolin (57.14 %), Cefuroxime and Cefuroxime Axetil(53.6 %) and Cefpodoxime (48.27 %). on the other hand, majority of bacterial isolates were susceptible to Levofloxacin, Tobramycin, Ciprofloxacin and Gentamycin with overall resistance rates of 0 %, 0 %, 6.6 %, and 11.11 %, respectively (Figure 1).

86.67% of the isolated bacteria were found to be resistant to one and more of the commonly used antibiotics listed in table-5. Among the total *Klebsiella* spp. Isolated all (100 %) of the isolates have developed resistant to one or more antibiotics used, among the total *Proteus* spp. isolated, the majority (77.8 %) of the isolates have developed resistance to one and more antibiotics used. Similarly, about three-fourth of (75 %) *Pseudomonas* spp. isolates were able to resist one and more antibiotics commonly used to treat them. All other bacterial isolates have also showed overall antibiotic resistance to one and more antibiotics used except *L.decarboxylata* and *C.koseri* which shows 100% sensitivity.

Discussion

ENT infection is a more frequent treatable health care problem worldwide, yet if left untreated; it can cause a serious complication such as a speech disorder, pain in patients and their family quality of life, and economic burden on the health care system [9]. The burden and prevalence of ENT infection are more intense in developing countries due to the poor living standard and hygienic conditions along with lack of proper nutrition [10]. Kumar [11] reported that from gram negative bacteria spp., *Proteus* spp. where responsible for most cases of ENT infections, Whereas Edwin [9] find out that *Pseudomonas* spp., *Proteus* spp. and *Klebsiella* spp., are the common bacteria that cause ENT infection in Japura India, similarly Argaw-Denboba [10] find out that *P. aeruginosa*, *Klebsiella* spp. *E. coli* are the bacteria

associated with ENT infection and *Pseudomonas aeruginosa* was the most prevalent aetiological agent of ENT infection in Benin city.

In this study, the main pathogenic bacterium associated with ENT infection was *Proteus mirabilis*. Followed by *Klebsiella pneumonia*, *Kelebsiella oxytoca* and *Pseudomonas aeruginosa*. respectively. Similarly, previously published articles from our study area and other parts of Ethiopia also reported *Proteus* spp. were the foremost bacteria associated with middle ear infection [9,10]. Although it needs a further nationwide study, taking into account our study finding and others, it seems that *Proteus Mirabilis* re the leading bacterial isolates associated with ENT infection in Ethiopia. Conversely, several other published data pieces from Africa and elsewhere in the world reported *Pseudomonas* spp., mainly *P. aeruginosa*, is the primary pathogenic bacteria associated with middle ear infection [12]. One possible explanation for this difference might be due to climate, poor living standard, hygienic condition and geographical variations between Ethiopia and other countries.

In general, Ampicillin, Cephalothin, Cefazolin, Cefuroxime and Cefuroxime had shown the highest antibiotic resistance rates to all bacterial pathogens isolated from ENT discharge, respectively. Correspondingly, Tetracycline, were the most clinically used antibiotics that showed a higher resistance rate to *Proteus* spp., which is in line with earlier reports

Table 2: Gram negative Bacteria Associated with ENT Infection in Arsho Advanced Medical Laboratory.

Isolates	Ear	Nose	Throat	Eye	Total (%)
<i>Pseudomonas aeruginosa</i>	3	-	-	-	3 (10 %)
<i>Proteus mirabilis</i>	7	2	-	-	9 (30 %)
<i>Kelebsiella oxytoca</i>	2	1	-	-	3 (10 %)
<i>Entrobacter cloacae</i>	2	-	-	-	2 (6.67 %)
<i>Citrobacter koseri</i>	1	-	-	-	1 (3.33 %)
<i>Citrobacter pauculus</i>	1	-	-	-	1 (3.33 %)
<i>Pseudomonas fulorescens</i>	1	-	-	-	1(3.33 %)
<i>Klebsiella pneumoniae</i>	1	5	-	-	6 (20 %)
<i>Acinetobacter baumannii</i>	1	-	-	-	1(3.33 %)
<i>E.coli</i>	1	-	-	-	1(3.33 %)
<i>Leclerciaa decarboxylate</i>	-	1	-	-	1(3.33 %)
<i>Proteus vulgaris</i>	-	1	-	-	1(3.33 %)

Table 3: Age Related Prevalence of gram negative bacterial Isolates in ENT Infection in Arsho Advanced Medical Laboratory.

Isolates	Age					
	<1	1-14	15-24	25-44	45-64	>65
<i>Pseudomonas aeruginosa</i>	-	-	2	1	-	-
<i>Proteus mirabilis</i>	-	4	3	1	1	-
<i>Kelebsiella oxytoca</i>	-	-	2	1	-	-
<i>Entrobacter cloacae</i>	-	-	-	2	-	-
<i>Citrobacter koseri</i>	-	-	-	-	1	-
<i>Citrobacter pauculus</i>	-	-	1	-	-	-
<i>Pseudomonas fulorescens</i>	-	-	1	-	-	-
<i>Klebsiella pneumoniae</i>	-	-	2	3	1	-
<i>Acinetobacter baumannii</i>	-	1	-	-	-	-
<i>E.coli</i>	-	1	-	-	-	-
<i>Leclerciaa decarboxylate</i>	-	1	-	-	-	-
<i>Proteus vulgaris</i>	-	-	1	-	-	-
Total	0	7	12	8	3	0
Percent (%)	0	23.33	40	26.67	10	0




Figure 1: Antibiotic susceptibility profiles of isolated bacteria from ENT.

Table 4: Overall antibiotic susceptibility profiles of isolated bacteria from ear discharge.

Type of antibiotics used	Frequency of Each antibiotic tested	Susceptibility patterns		
		Resistant	Sensitive	Intermediate
Ampicillin	26	80.8 % (21)	15.4 % (4)	3.8 % (1)
Augmentin	28	25 % (7)	75 % (21)	-
Cefazolin	28	57.14 % (16)	39.3 % (11)	3.6 % (1)
Cefepime	30	26.7 % (8)	73.3 % (22)	-
Cefoxitin	29	44.83 % (13)	48.27 % (14)	6.9 % (2)
Cefpodoxime	29	48.27 % (14)	51.73 % (15)	-
Ceftazidime	30	30 % (9)	70 % (21)	-
Ceftriaxone	30	36.67 % (11)	63.33 % (19)	-
Cefuroxime	28	53.6 % (15)	39.3 % (11)	7.14 % (2)
Cefuroxime Axetil	28	53.6 % (15)	46.4 % (13)	7.14 % (2)
Cephalothin	29	58.6 % (17)	37.9 % (11)	3.4 % (1)
Ciprofloxacin	30	6.67 % (2)	90 % (27)	3.33 % (1)
Co-trimoxazole	27	29.63 % (8)	70.37 % (19)	-
Gentamycin	27	11.11 % (3)	85.2 % (23)	3.7 % (1)
Levofloxacin	29	-	93.1 % (27)	6.9 % (2)
Piperacillin/Tazobactam	28	28.6 % (8)	71.4 % (20)	-
Tetracycline	28	42.9 % (12)	57.1 % (16)	-
Tobramycin	28	-	96.4 % (27)	3.6 % (1)

from Ethiopia, Nigeria, and Egypt [13]. Among the total *Klebsiella* spp. and *Proteus* spp. isolated during the study period about 100% and 77.8% of the isolated bacteria have developed resistance to one and more antibiotics that were once in clinical use, respectively. Overall, more than 60% of the bacterial isolates of this study were characterized as multi-antibiotic-resistant pathogenic bacteria. The reason for this high degree of multiantibiotic resistance might be connected to inappropriate use of antibiotics, including incomplete dose, self-medication, and poor infection prevention and control practices as indicated by the recent WHO antimicrobial resistance report. An Interestingly, our study revealed almost all the isolated pathogenic bacteria were considerably susceptible to Levofloxacin, Tobramycin, Ciprofloxacin and Gentamicin. Particularly, Levofloxacin, and Tobramycin were shown to be highly effective for pathogenic bacteria associated with ENT

infection in this study. Likewise, with a slight variation several other authors have shown a similar high efficiency of ciprofloxacin against these bacterial species. Thus, we propose that Levofloxacin, and Tobramycin can be taken as a first-line optional treatment for ENT infection attending in Arsho Advanced medical Laboratory, Addis Ababa, Ethiopia. This suggestion holds true, since the current first-line treatment for both acute and chronic otitis media in Ethiopia is Ampicillin and Augmentin which already showed a high resistance rate (80.8 % and 25 % respectively) for the majority of the bacterial isolates in this study.

Conclusion and Recommendation

This study showed that the risk of acquiring particularly ear infection is strongly associated in *Proteus mirabilis*, *Pseudomonas aeruginosa* and *Klebsiella oxytoca* were the three predominant bacteria isolates from patient ear

Table 5: Antibiotic susceptibility test of pathogenic bacteria isolated from ENT patients.

Organisms	AMP	AMC	TZP	CFA	CFZ	CFU	CFXA	FOX	CPD	CAZ	CRO	CFP	GM	TBM	CIP	LEV	TEC	SXT
<i>P.mirabilis</i>	-		S	S	-		S	S	S	S	S	S		S	S	S	S	
<i>P.mirabilis</i>	R	S	R	R	R	R	R	R	R	R	R	R	S	S	S	S	R	R
<i>P.mirabilis</i>	R	S	R	R	R	R	R	R	R	R	R	R	R	S	R	S	R	R
<i>P.mirabilis</i>	R	S	R	R	R	R	R	R	R	R	R	R	I	S	S	S	R	R
<i>P.mirabilis</i>	-		-		-	-	S		-	S	S	S			S		-	
<i>P.mirabilis</i>	S	S	S	R	S	S	S	S	S	S	S	S	S	S	S	S	R	S
<i>P.mirabilis</i>	R	S	R	R	R	R	R	S	R	R	R	R	R	S	S	S	R	R
<i>P.mirabilis</i>	R	S	R	R	R	R	R	I	R	R	R	R	S	S	S	S	R	R
<i>P.vulgaris</i>	R	S	R	R	R	R	R	R	R	R	R	R	S	S	S	S	R	S
<i>P.aeruginosa</i>	R	R	R	R	R	R	R	R	R	R	R	S	S	S	S	S	R	R
<i>P.aeruginosa</i>	R	R	S	R	R	R	R	R	R	S	R	S	S	S	I	I	R	R
<i>P.aeruginosa</i>	R	R	S	R	R	R	R	R	R	S	R	S	S	S	S	S	R	R
<i>P. fluorescens</i>	S	S	S	S	S	S	S	S	S	S	S	S	S	S	S	S	S	S
<i>K.oxytoca</i>	R	S	S	I	I	R	R	S	S	S	S	S	S	S	S	S	S	S
<i>K.oxytoca</i>	R	R	S	S	R	S	S	S	S	S	S	S	S	S	S	S	S	S
<i>K.oxytoca</i>	R	S	S	S	S	R	S	S	S	S	S	S	S	S	S	S	S	S
<i>K.pneumoneae</i>	R	S	S	S	S	S	S	S	S	S	S	S	S	S	S	S	S	S
<i>K.pneumoneae</i>	R	S	S	S	S	S	S	S	S	S	S	S	S	S	S	S	S	S
<i>K.pneumoneae</i>	R	S	S	S	S	S	S	S	S	S	S	S	S	S	S	S	S	S
<i>K.pneumoneae</i>	R	S	S	S	S	S	S	S	S	S	S	S	S	S	S	S	S	S
<i>K.pneumoneae</i>	R	S	S	R	S	R	S	R	R	S	S	S	S	S	S	S	S	S
<i>K.pneumoneae</i>	R	S	S	R	R	S	I	R	S	S	S	S	S	S	S	S	S	S
<i>E.cloacae</i>	R	R	S	R	R	I	R	R	R	S	S	S	S	S	S	S	S	S
<i>E.cloacae</i>		R	S	R	R	I	R	R	R	S	S	S	S	S	S	S	R	S
<i>C.koseri</i>		S	S	S	S	S	I	S	S	S	S	S	S	S	S	S	S	S
<i>C.pauclus</i>	S	S		R	R	S	S	S	S	S	S				S			
<i>A.baumannii</i>	I	S	S	S	S	R	R	R	S	S	S	S	S	S	S	S	S	S
<i>E.coli</i>	R	R	S	R	R	R	R	I	R	R	R	R	R	I	R	I	S	S
<i>L. adecarboxylata</i>	S	S	S	S	S	S	S	S	S	S	S	S	S	S	S	S	S	S

discharges suspected of otitis media whereas, in nose infection *Klebsiella pneumonia* was the most predominant isolated organism. Almost all the isolated bacteria showed a considerable level of resistance to one antibiotic that are commonly used in health institutes; particularly, majority of isolated bacteria were found to be highly resistant to Ampicillin Cephalothin, Cefazolin, Cefuroxime and Cefuroxime Axetil and Cefpodoxime treatments. Among the total *Klebsiella* spp. Isolated all (100 %) of the isolates have developed resistant to one or more antibiotics used, among the total *Proteus* spp. isolated, the majority (77.8 %) of the isolates have developed resistance to one and more antibiotics used. This study also indicated Levofloxacin, and Tobramycin are effective against all the bacterial isolates and most was highly sensitive to Tobramycin. In general, the result of this study revealed that antibiotic-resistant bacteria are alarmingly increasing in Arsho Advanced Medical Laboratory study area, Addis Ababa, Ethiopia, and becoming a major public health problem in patients with ENT infection. Therefore, we recommend nationwide antimicrobial inspection to make the right recommendation of alternative antibiotics along with strict devotion to antibiotic policy to reduce the spread of drug resistant microbes in the country.

Acknowledgment

We would like to thank all staff members of Arsho Advanced Medical Laboratory, microbiology department.

References

1. Witsell DL, Dolor RJ, Boile JN, Stinnet SS (2001) Exploring health related quality of life in patients with diseases of the ear, nose and throat; A multicenter observation study. *Otolaryngology Head and Neck Survey* 125: 288-298.
2. Albert PW (1999) Pediatric ear, nose and throat service demands and resources: a global perspective. *International Journal of Pediatrics Otolaryngology* 18: 136-139.
3. Chibuike I, Reginald AO, Solomon UC, Ifeanyi AO, Conrad J, et al. (2013) Prevalence and Antibiotic Susceptibility Patterns of Methicillin-Resistant *Staphylococcus Aureus* (MRSA) Isolated from Healthy Inhabitants of Uturu Rural Communities, Abia State, Nigeria. *Journal of Natural Science Research* 3: 85-91.
4. Obiajurn IOC, Chukuezi AB (2013) Microbiological assessment of ENT Infections and disease: Case study at Orlu, Imo state, Nigeria. *Journal of Research in Nursing and Midwifery* 2: 72-76.
5. Newton EV, Macharia J, Muswe P, Ototo B, Kan SW (2001) Evaluation of the use of a questionnaire to detect hearing loss in Kenyan pre-school children. *International Journal of Pediatric Otolaryngology* 57: 229-234.
6. Rover MM, Dekok IMCM, Schilder AGM (2006) Risk factors for Otitis media: An International perspective. *International Journal of Pediatric Otolaryngology* 2: 1251-1256.
7. Noh KT, Kim CS (1985) The changing pattern of otitis media in Korea. *Int J Pediatric Otorhinolaryngol* 9: 77-87.
8. Chibuike I, Reginald AO, Solomon UC, Ifeanyi AO, Conrad J, et al. (2013) Prevalence and Antibiotic Susceptibility Patterns of Methicillin Resistant *Staphylococcus Aureus* (MRSA) Isolated from Healthy Inhabitants of Uturu Rural Communities, Abia State, Nigeria. *Journal of Natural Science Research* 3: 85-91.

9. Edwin B, Prasanna V, Kannan I, Katiyar VH, Dhanapal E (2014) Incidence of bacterial colonization in the oropharynx of patients with ear, nose and throat infections. *International Journal of Medical Science and Public Health* 3: 931-934.
10. Argaw-Denboba A, Agalu Abejew A, Gashaw Mekonnen A (2015) Antibiotic-Resistant Bacteria Are Major Threats of Otitis Media in Wollo Area, Northeastern Ethiopia. *International Journal of Microbiology*.
11. Kumar RP, Sriya steva M, Sharma S, Rishi PS, Nwawan P (2013) Isolation and Antimicrobial Sensitivity profile of Bacterial Agents in Chronic Supportive Otitis media in patients at NIMS Hospital, Jaipur. *Journal of Research in Nursing and Midwifery* 2: 265-269.
12. Afolabi OA, Salaudeen AG, Ologe FE, Nwabuisi C, Nwawolo CC (2012) Pattern of bacterial isolates in the middle ear discharge of patients with chronic suppurative otitis media in a tertiary hospital in north central Nigeria. *African Health Sciences* 12: 362-368.
13. Seid A, Deribe F, Ali K, Kibru G (2013) Bacterial otitis media in all age group of patients seen at Dessie referral hospital, North East Ethiopia. *Egyptian Journal of Ear, Nose, Throat and Allied Sciences* 14: 73-78.