

Systematic Review on the Efficacy of Interventions for Fear of Childbirth, Anxiety and Fear in Pregnant Women

Birner Carmen*

Grosse Gerlind

*Department of Social and Education Sciences, Potsdam University of Applied Sciences,
Germany*

Abstract

Background: Fears and anxieties during pregnancy and childbirth are a frequent phenomenon and can have negative consequences on wellbeing, psychological health and birth outcomes. Therefore, it is important to focus on the interventions to reduce those fears and anxieties during pregnancy and childbirth. A systematic review was conducted to examine the current literature on psychological interventions to reduce anxieties and fears during pregnancy and childbirth. Scopus and PubMed were searched from 2015 up until December 2020 for relevant studies. Included were pregnant women, with no restriction on age ranges or parity. Entered in the review were quantitative studies, including randomized controlled trials (RCTs), non-randomized controlled trials as well as treatment evaluations. After reviewing titles, abstracts and studies, 72 studies were included in this review as they met the inclusion criteria. Standard methodological procedures for systematic reviews were used. The quality assessment of included articles was done by using the Quality Assessment Tool for Quantitative Studies (EPHPP).

Results: The main results of this review concern the fear and anxiety reducing effects of psychoeducation, relaxation techniques, guided imagery, supportive care through a midwife, group discussion, "lifestyle based education", writing therapy, cognitive behavioral therapy groups and stress intervention, individual structured psychotherapy, communication skills training, counselling approaches (except distraction techniques), a motivational interviewing psychotherapy, emotional freedom techniques, breathing awareness and different hypnotherapeutic techniques on different fears and anxieties during pregnancy and childbirth. For mindfulness-based interventions mixed results are found. The effect of an acceptance and commitment therapy, biofeedback interventions, a mind body intervention, mental health training courses, the group intervention Nyytti® as well as cognitive analytic therapy is unclear, due to weak study ratings. Antenatal class attendance reduced delivery fear significantly only in first time mothers. An internet-based problem-solving treatment did not reduce anxiety during pregnancy.

Conclusion: A broad range of interventions show positive effects on fear of childbirth and fear and anxiety in pregnancy. Further research should address other acknowledged psychotherapeutic practices, like psychodynamic as well as systemic interventions, as they are underrepresented within this review. Furthermore, there is a need for manualized therapeutic interventions, with regards to a combination of effective intervention components.

Article Information

Article Type: Research

Article Number: JPI-144

Received Date: 10 May, 2021

Accepted Date: 22 September, 2021

Published Date: 29 September, 2021

***Corresponding author:** Birner Carmen, Department of Social and Education Sciences, Potsdam University of Applied Sciences, Kiepenheuerallee 5, 14469 Potsdam, Germany. Tel: +4933158000; E-mail: carmen.birner@fh-potsdam.de

Citation: Carmen B, Gerlind G (2021) Systematic Review on the Efficacy of Interventions for Fear of Childbirth, Anxiety and Fear in Pregnant Women. J Pediat Infants Vol: 4, Issu: 2 (66-90).

Copyright: © 2021 Carmen B et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Keywords: Fear, Anxiety, Childbirth, Pregnancy, Intervention.

Introduction

In the current literature, the prevalence rate of high levels of fear of childbirth is stated as 36.7% in Ireland and in India the prevalence rate of severe pregnancy anxiety levels reached up to 22% [1,2].

Women with very high scores on Fear of Childbirth (FOC) or Tocophobia often suffer under longer birth processes and stronger to unbearable pain compared to women with less fear [2-8]. FOC is reported as one of the most common reasons for unnecessary cesarean sections [2-8]. Compared to women with low levels of FOC, women with intermediate or high levels of FOC seem to have more negative birth experiences [9]. FOC can not only have a negative impact on the birth process, but also influences the wellbeing during pregnancy [10].

There is also evidence for a connection between FOC, postpartal depressions and traumatic stress symptoms [11-14]. Furthermore, severe FOC or even anxiety may result in pre-term delivery, bonding issues and behavioral/emotional problems of the infant [15-18].

Besides those findings, the current corona pandemic and a resulting increased fear of COVID-19 is a predictor for worries in pregnant women [19]. As possible worries the authors state the “worry about fetus health or mothers’ own health and worry going to hospital” [19]. Molgora and Accordini also stated in their study, that the time of the pandemic has a significant negative impact on the pregnant women’s wellbeing [20].

A number of systematic reviews regarding interventions to treat fear of childbirth exist in the literature of the last five years. There are general systematic reviews and meta-analyses, that list different interventions and their effect on “fear of childbirth” [21-23]. Further systematic reviews focus on pregnancy specific anxiety, as well a mental disorders during pregnancy [24-26].

Besides more general reviews, systematic reviews specifically focussing on mindfulness interventions, psychotherapy interventions, e-health and technology based interventions exist [27-39]. Further meta-analyses focus on the effect of expressive writing, psychoeducation interventions and hypnosis based interventions on anxiety related to pregnancy [40-42].

The reviews of Bright et al., Brixval et al. and O’Connell et al. are not listed as only study protocols were found [43-45]. The mentioned reviews included studies up to the year 2019.

The present systematic literature review is the broadest and up to December 2020 most recent overview regarding the effects of psychological interventions on fear and anxiety related to pregnancy and childbirth published in the last five years with 72 included and rated studies. While some of the past reviews focused only on certain outcome variables (e.g. only fear of childbirth as a narrow topic), this present

review focusses on broader fears and anxieties regarding the whole pregnancy and childbirth process and therefore addresses a research gap [21,23,46]. This systematic review encloses studies up until December 2020.

The aim of this systematic review was to examine the effect of psychological interventions on “fear of childbirth” as well as fears and anxieties during pregnancy.

Previous reviews stated positive effects of psychological interventions. Hypnosis based, psychotherapeutic interventions and psychoeducation seem to have a positive impact on fear of childbirth [25,42,47]. There is a need to keep those findings updated and an existing research gap to review further interventions stated within the literature.

Definitions

There is no clearly delimitable and common definition of fear of childbirth (FOC) in the literature. It is also difficult to draw a line between subclinical, phobic and pathological levels of FOC [48]. To give an overview over existing terms in the literature, this paper makes an attempt to define different phrases related to the term FOC. This systematic review focusses besides FOC on different anxieties and fears during pregnancy and childbirth.

FOC (Fear of childbirth)

Areskog defined „fear of childbirth“ (FOC) first in a population of Swedish pregnant women as: “a strong anxiety which had impaired their [the women’s] daily functioning and wellbeing”. Later, during the 2000s, a study from Sweden defined FOC as belonging to “the family of anxiety disorders” [49,50].

Klabbers pointed out, that FOC is an anxiety disorder or phobic fear [51].

In the classification of diseases – 10 (ICD 10) fear of childbirth could most likely be listed under code O99.8 as “other specific diseases and conditions complicating pregnancy, childbirth, or puerperium” [52].

Wijma describes “clinical FOC”, as a “disabling fear that interferes with occupational or academic functioning, with domestic and social activities or with relationships”. The symptoms of FOC could be characterised as “worries or extreme fear” [53,54].

FOC has different manifestations. It is “assumed to be a continuum with no or low fear on one end, and severe or extreme fear on the other” and is clinically relevant “if it affects a woman’s quality of life” [23].

FOC can also be classified into primary FOC, which occurs in nulliparous women and secondary FOC relating to women who already had traumatic birth experiences [48,55]. A third form is FOC as a symptom of prenatal depression [51,55,56].

Tocophobia

Primary tocophobia is defined as “severe fear precedes conception and leads to avoidance of tokos (Greek: childbirth)”, while secondary tocophobia is a “phobic fear resulting from a distressing or even traumatising childbirth experience” [23]. It is characterised as an “unreasoning dread

of childbirth” relating to women in a “specific and harrowing condition” including a “pathological dread” and “avoidance of childbirth” [56]. Tocophobia is “a specific anxiety or fear of death during parturition precedes pregnancy” that is “so intense that tokos (childbirth) is avoided whenever possible; this is a phobic state called tocophobia” [57].

Bhatia and Jhanjee defined tocophobia as “a pathological fear of pregnancy” and indicated the pathological aspect of tocophobia, which can result in avoidance of childbirth [54,58]. The authors distinguish tocophobia – similar to the classification of FOC - between primary fear of childbirth, in women without previous pregnancy experience and secondary fear of childbirth related to a “traumatic obstetric event in previous pregnancy” [54].

Tocophobia often is defined by $\geq 85/165$ points on the Assessments Wijma Delivery Expectancy Questionnaire (W-DEQ A) [8,59].

A recent systematic review determined tocophobia to be synonymous with severe FOC [59]. Based on this conclusion this present paper also refers to severe/high FOC as synonymous to tocophobia.

Childbirth anxiety (CA)

Wijma and Wijma define childbirth anxiety as follows: “When a woman is afraid of the situation where a child will or is to be born [...] CA covers the whole continuum from a little fear that is easy to cope with to phobic fear, when the woman wants to avoid the situation by all means”[60].

Perinatal anxiety (PNA) and perinatal generalized anxiety disorder (GAD)

Harrison, Moore and Lazard characterized the term perinatal anxiety and Misri et al. introduced the term Perinatal Generalized Anxiety Disorder (GAD), which is defined as “excessive, uncontrollable worry that can cause functional impairment” [61,62].

Further forms of pregnant related anxieties

Pregnancy can be accompanied by a variety of anxiety disorders, like panic, disorder with or without agoraphobia, obsessive-compulsive disorder, generalized anxiety disorder, specific phobia, social anxiety disorder and post-traumatic stress disorder [18].

Methods

Criteria for considering studies

Papers included in this systematic review were limited to publications in English and German language only with the restriction for publication year between 2015 and December 2020.

Inclusion criteria: The inclusion criteria for this systematic review were outcomes regarding fears or anxieties during pregnancy and childbirth. Different definitions of the concept “fear of childbirth” and the understanding of fears and anxieties during childbirth were admitted. Besides, varying outcome measurements were valid. Pregnant women (primi-/nulli- and/or multiparous as well as primi-/nulli- and/or multigravida) with no restriction to age

ranges were included. The interventions were restricted to psychological interventions, biofeedback interventions, mindfulness-based interventions and midwife counselling studies. Included studies focused on the prenatal period. Only intervention studies were included (no correlation studies about personality traits).

Exclusion criteria: Excluded were study protocols, qualitative studies, reviews, case series designs, case reports, consensus bundles, medical research counsel frameworks, studies with no described study design and uncorrected proof studies. Studies of pregnant women with specific somatic complains, (in)fertility or abortion studies, yoga interventions, pharmacological interventions (including psychopharmacology), music interventions, spiritual interventions, art therapy as only intervention, and aroma therapy studies were excluded. Not included were medical studies, studies on pregnancy loss or sleeping problems, postpartum studies, traumatic birth studies, studies focusing on depression only, sport or physical activity interventions, studies relating to stillbirth, and ultrasound interventions. Studies written in other languages than German or English were also excluded.

Search methods for identification of studies

The electronic databases PubMed and Scopus were searched for articles using the terms “fear”, “anxiety”, “pregnancy”, “childbirth”, “intervention” from 2015 up to December 2020.

The initial search yielded a total of 3029 studies, after setting the time (year 2015-2020) and language filters a total of 2027 studies were displayed and a total of 1426 studies were screened for this review, after removing all duplicates. Further studies were excluded as they were either not relevant to the review or did not meet the inclusion criteria or were not found. 72 records were screened. See figure 1 for the summary of search item identification. For the final included studies and their results see table 1.

Data collection and analysis

One person was included in the data collection, management and analysis of the studies. No software tools were used to support selection of studies. With an excel programme duplicates were analyzed. No standardized data collection forms were used. The data items are described in table 1.

Quality assessment and risk of bias in included studies

Of the 72 studies included, 22, 31 and 19, respectively, received a strong, moderate and weak rating on the “Quality Assessment Tool for Quantitative Studies” of the “Effective Public Health Practice Project (EPHPP)”[144]. The risk of bias in the included studies was assessed with this tool. One author was involved in the assessment of risk of bias in included studies. All studies (strong, moderate, weak ratings) were included in the analysis and interpretation. The ratings are listed in table 2.




Figure 1: Systematic Review Profile based on the prisma flow diagram [141].

Dealing with missing data

Few studies without access were excluded from analysis: Nasiri et al., Kao et al., Jahdi et al., Anton and David, Soltani et al., Najafi et al., Hennelly et al. [63-69]. No authors or sponsors were contacted to obtain missing information or clarify the information available. Missing data (e.g. the period of time of data collection) within the viewed studies were marked as such in table 1.

Results

Description of Studies

Characteristics of the included studies

Details of the search results are presented in table 1. 72 studies fulfilled the inclusion criteria, with a total of 8288 pregnant women. The 72 studies were the basis of the findings within this review. The studies were conducted across 18 countries with most studies from Iran (32 studies), second most from Turkey (7 studies) and third most from Sweden, Netherlands and UK (4 studies), China, Finland and USA (3 each), India, Australia and France (2 each), Germany, Iceland, Malawi, Poland, Japan and Malaysia (1 each). Most studies were RCTs, with 47 RCT study designs. The period of time of data collection ranged from 2007 up until 2020.

10 studies included primi- and multiparous women, 7 studies included nulli- and multiparous women, 1 study included nulligravidae and 5 studies primigravidae women. 9 studies included primiparous and 9 nulliparous women. 3

studies included primi- and multigravidae women. 1 study included primi antenatal women. For 27 studies parity or gravidity could not be stated.

Outcome variables

14 studies focused on „fear of childbirth” as an outcome [47,48,70-92].

Some studies focused on “state/trait anxiety during pregnancy”, while others focused only on “state anxiety during pregnancy [92-107].

Some studies had “pregnancy related anxiety as an outcome” and others focused on “anxiety during pregnancy” [96,108-125].

Another outcome was “general anxiety during pregnancy” [126-130].

Few studies focused on specific outcome variables like “mental health of pregnant women – anxiety”, “perinatal mood and anxiety disorders”, “anxiety of pregnant women undergoing interventional prenatal diagnosis”, “labor fear”, “fear of women undergoing labor”, “pain catastrophizing”, “pregnancy worries and stress”, “fear of delivery” and “perceived stress during pregnancy” [131-139].

Interventions applied to pregnant women

Online vs. offline: Interventions were delivered in two forms: Via the internet respectively online or digital or offline respectively face to face [81,83,87,89,91,95,108,118,126-

First Autor, Year	Country/ date of data collection/ study design/ EPHPP rating	Target population (sample size/age (mean, ± SD)/ gravidity and parity)	Intervention/ comparators	Study outcome: Interventions Key findings	Measurements (for fear and anxiety during childbirth and FOC)	For this study relevant research topic
Aksoy Derya (2021)	Turkey 2020 RCT moderate	<i>N</i> =96 Age: IG: 28.70 ± 4.73 CG: 28.06 ± 4.12 Not stated	IG: individual tele-education (interactive education and consultancy provided by phone calls, text message and digital education booklet) CG: No intervention	The posttest PRAQ-R2 total mean scores ($t=-4.095$, $p=.000$) of the pregnant women in the IG and CG, as well as the subscales “fear of giving birth” ($t=-3.275$, $p=.001$) and “worries of bearing a physically or mentally handicapped child” ($t=-4.354$, $p=.000$) showed a statistically significant difference between the groups. The subscale “concerns about own appearance” did not show a statistical difference between the groups. When the intragroup comparisons of the pre- and posttest in the IG were examined, their “pretest prenatal distress”, “fear of giving birth”, “worries of bearing a physically or mentally handicapped child” and “pregnancy-related anxiety” total mean scores were significantly lower than their posttest mean scores ($p<.05$). In the CG only the “pretest fear of giving birth” subscale mean score was significantly lower than the posttest mean score ($p<.05$).	Pregnancy Related Anxiety Questionnaire-Revised-2 (PRAQ-R2) Revised Prenatal Distress Questionnaire (NuPDQ)	Preg-nancy related anxiety
Montazeri (2020)	Iran 2018 RCT moderate	<i>N</i> = 70 Age: IG: 27.5 ± 5.9 CG: 27.7 ± 5.8 Not stated	IG: Three protocol-based writing therapy sessions CG: routine pregnancy care	The results of the independent <i>t</i> -test showed no significant difference in the mean score of pre-intervention anxiety in the IG and CG ($p=.287$). According to ANCOVA with baseline score adjustment, the score of anxiety had a significant reduction in the IG compared to the CG (adjusted mean difference: -6.8; 95% confidence interval: -9.1 to -4.5; $p<.001$).	Beck anxiety inventory	Anxiety during pregnancy
Waters (2020)	UK Date of data collection not stated open-label pilot study weak	<i>N</i> = 74 Age: 33.5 (3.87) Primi- and Multiparous	8-week, group-delivered Acceptance and Commitment Therapy (ACT) intervention	At post-treatment, 38 of 55 women (69%) demonstrated a statistically reliable decrease in global distress ($d=0.99$).	Diagnoses of moderate-to-severe anxiety disorders were made by the PCMHS team Consultant Perinatal Psychiatrist (SS) (reviewed all routinely obtained clinical data against ICD-10 criteria). Clinical Outcomes in Routine Evaluation – Outcome Measure (CORE-OM)	Perinatal mood and anxiety disorders
Zarenejad (2020)	Iran Date of data collection not stated RCT strong	<i>N</i> =70 Age: IG: 27 ± 5 CG: 24.5 ± 5 0 Not stated	IG: received 6 mindfulness-based stress reduction (MBSR) training sessions CG: routine care	The results of analysis of variance with repeated measures in assessing the changes in pregnancy anxiety score before, immediately after, and 1 month after the intervention showed that the length of time affects the anxiety score of pregnancy by decreasing it ($p=.03$) and that a significant difference was observed between the two groups in this regard ($p=.001$). After the intervention, the CG showed significant higher scores in anxiety compared to the IG.	Pregnancy-Related Anxiety Questionnaire	Preg-nancy related anxiety
Firouzan (2020)	Iran 2019 RCT strong	<i>N</i> =80 Age: IG: 26.27 ± 4.48 CG: 25.87 ± 4.58 Nulligravida	IG: face-to-face counselling sessions based on the BELIEF protocol + telephone-counselling sessions CG: prenatal routine care	After adjusting for the pretest scores, there was a significant difference between the IG and CG on post-test scores of W-DEQ-A ($F(1,65)=100.42$, $p=.0001$, partial eta squared = .60). The IG got lower scores on W-DEQ-A at post-test than the CG, indicating that the BELIEF protocol was effective in decreasing childbirth fear.	W-DEQ A	FOC
Kang (2020)	China 2012-2014 RCT weak	<i>N</i> =100 Age: 26.9 ± 1.5 Not stated	IG: psychological intervention CG: Routine Nursing Care	Postoperative SAS scores were significantly lower in the IG than in the CG and the differences were statistically significant ($p < 0.01$). In the CG, differences in anxiety and fear levels were not statistically significant between preoperation and postoperation ($p > 0.05$).	Self-rating Anxiety Scale (SAS)	Anxiety of pregnant women under-going interventional prenatal diagnosis

Alipour (2020)	Iran 2017-2018 RCT moderate	<i>N</i> =54 Age: IG: 29.1 (4.3) CG: 29.4 (4.5) Primi- and multiparous	IG: communication skills training package + couple-based intervention CG: two sessions of childbirth preparation + after the completion of the third phase of the study: given educational pamphlets	The level of anxiety three months after intervention was lower ($p=.001$) in the IG than in the CG. The results showed the impact of group in the level of anxiety ($p<.001$) was significant. During the study follow-ups in the IG, a significant change in the level of anxiety ($p<.001$) occurred.	Questions related to the subscales of depression and anxiety of General Health Questionnaire (GHQ)	Anxiety during pregnancy
Goetz (2020)	Germany 2019 prospective pilot study with an explorative study design weak	<i>N</i> =68 Age: 32.07 (4.74) Not stated	Intervention: electronic Mindfulness-based interventions (eMBIs)	After completing the 1-week electronic course on mindfulness, the participants showed a significant reduction in the mean state anxiety levels ($p<.05$).	State-Trait Anxiety Inventory (STAI-S) Pregnancy-Related Anxiety Questionnaire (PRAQ-R)	Preg-nancy related anxiety State/ Trait Anxiety
Sridhar (2020)	USA 2018 Pilot feasibility study moderate	<i>N</i> =30 Age: 30.1 (7.4) Not stated	IG: Participants could choose any of the three available virtual reality (VR) environments (dream beach, Iceland, dolphins) CG: receiving standard care	The median decrease in the VAS anxiety score from before to after the procedure was greater in the IG than in the CG (Wilcoxon rank-sum, $p=.3324$) All but one participant reported that VR was either very effective (53%) or somewhat effective (40%) in relieving anxiety during and after the procedure.	Modified Amsterdam Preoperative Anxiety and Information Scale (APAIS) + a visual analogue scale (VAS) for anxiety, ranging from 0 (minimum anxiety) to 10 (maximum anxiety)	Anxiety during pregnancy
Esfandiari (2020)	Iran 2018-2019 RCT moderate	<i>N</i> =80 Age: IG: 27.87(5.26) CG: 23.72(4.27) Not stated	IG: group supportive counseling (SC) CG: antenatal usual care (AUC)	In the IG scores of state-anxiety were reduced more remarkably than in the CG with a large effect size ($B=-8.47$, $p= <0.001$, $\eta^2 = 0.40$).	Spielberger State-Anxiety Inventory (STAI-Y)	State anxiety during pregnancy
Mirtabar (2020)	Iran 2017-2018 RCT strong	<i>N</i> =60 Age: 29.0 ± 5 Not stated	IG: received individual structured psychotherapy + preterm labor inpatient medical care CG: inpatient medical care for preterm labor	Both the IG and CG had significant reductions in the mean scores of state-anxiety and pregnancy distress from the baseline to end of study ($p<.05$). The ANCOVA tests determined that the IG had a significant improvement in the state-anxiety scores compared with the CG ($p<.001$).	State-Anxiety Inventory (STAI)	State anxiety in preterm labor Preg-nancy distress
Abbasi (2020)	Iran 2015-2016 RCT moderate	<i>N</i> =153 Age: IG Educational software :25.5 (3.8) IG: Educational Booklet: 25.9 (3.6) Control: 25.1 (3.2) Not stated	IG Educational software: studied the educational content of the educational software IG: Educational Booklet: studied the educational content of the educational booklet CG: routine care	The average state anxiety score in the educational software group and the educational booklet group was significantly lower than the CG ($p<.001$). Also, the mean state anxiety score in the educational software group was significantly decreased compared to the educational booklet group after the intervention ($p<.001$). The average score of trait anxiety in the educational software group and the educational booklet group was significantly lower than the control group ($p<.001$). Also, there was no significant difference between the two intervention groups ($p=.952$)	State-Anxiety Inventory (STAI)	State and trait anxiety during pregnancy
Larsson (in press)	Sweden 2014- 2015 RCT weak	<i>N</i> =258 Age (n, %) <25: 14 (10.4) 25-35: 100 (74.6) >35: 20 (14.9) Primi- and multiparous	IG: internet-based cognitive behavioral therapy (iCBT) CG: standard care (i.e. counseling with midwives)	No statistically significant difference in the perceptions of the birth experience, regardless of treatment method for fear of birth.	Fear of birth scale	FOC

Swift (in press)	Iceland 2017-2018 quasi-experimental controlled trial strong	$N=92$ Age: IG: 28.3 (5.1) CG: 27.9 (4.4) Not stated	IG: Enhanced Antenatal Care (EAC) CG: usual antenatal care	At baseline, a higher proportion of IG participants (28%) reported high fear (>60 points) compared with women in CG (21%). By T2 fewer women reported high fear of birth in IG (9.4%) compared with CG (15.0%). For the full sample, the mean childbirth fear change score was 7.2 points among women in IG and -3.0 points among women in usual care ($p=0.315$). Based on Cohen's criteria the effect of participating in IG on the reduction in mean childbirth fear was small (Cohen's $d=0.21$). Restricting the main analysis to women who had not attended classes alongside antenatal care ($n=26$) resulted in a large effect size difference in fear change between women in IG and CG (Cohen's $d=-0.83$), with a change score of -14.1 points among women in IG and a slight increase in fear among women in CG (1.2 points; $p=.003$).	Fear of birth scale (FOBS)	FOC
Shahsavani (2020)	Iran 2018 quasi-experimental study strong	$N=102$ Age: IG: 28.10 (± 5.20) CG: 28.69 (± 5.31) Nulliparous	IG: Internet-based guided self-help cognitive- behavioral therapy (I-GSH-CBT) CG: normal pregnancy care	The IG intervention could significantly reduce the scores of childbirth fear ($p=.002$). The fear scores in the CG were significantly increased in parallel with the IG intervention ($p<.001$).	W-DEQ	FOC
Boz (2020)	Turkey 2018 RCT moderate	$N=24$ Age: 28.21 (± 4.37) Nulliparous	IG: Psychoeducation Program based on Human Caring Theory in The Management of Fear of Childbirth CG: Antenatal education classes group	The FOC of women from pretest to posttest was statistically more reduced in the IG compared to the CG ($p=.000$).	W-DEQ-A/B	FOC
Abdollahi (2020)	Iran 2018 RCT moderate	$N=70$ Age (range): aged 18–50 Not stated	IG: Motivational Interviewing (MI) Psychotherapy CG: Prenatal usual care (PUC)	The total score of W-DEQ declined more considerably in the IG than in the CG between pre-trial (T0) and post-trial (T1), with a large effect size ($B=-23.54$, $p<.001$, $\eta^2 = 0.27$). Scores of the six subscales of W-DEQ diminished more substantially in psychotherapy than in prenatal usual care.	W-DEQ Spielberger state anxiety	FOC
Hamilton (2020)	UK Date of data collection could not be stated RCT weak	$N=39$ Age: TAU+CAT: 30.2 (6.4) TAU: 31 (2.9) Not stated	IG: cognitive analytic therapy (CAT) plus treatment as usual (TAU) CG: treatment as usual (TAU)	The analysis found no difference in the primary outcome. The STAI scale at 24 weeks after randomization between the groups, with an adjusted difference in means of 6.1 points (95% CI: -4.2 to 16.3) was in favor of CAT for the State domain and 6.2 points (95% CI: -2.8 to 15.2) for the Trait domain. The IG having lower (better) STAI scores at all four post-randomization assessment points than the CG. For the four post-randomization repeated STAI measures, a simple summary measure for each individual patient, the average post-randomization score was calculated. Average post-randomization STAI scores were compared between the two arms (CAT and TAU), again with analyses unadjusted and adjusted for covariates. All the 95% CIs for the difference in mean follow-up scores between the CAT and TAU groups, include zero, which is compatible with no difference in outcomes between the randomized groups.	Spielberger State/ Trait Anxiety Inventory (STAI)	State/ Trait Anxiety
Noorbala (2019)	Iran 2015–2018 Clinical Trial Study weak	$N=202$ Age: 27.92 \pm 5.41 Not stated	IG: life skills and stress management training, supportive psychotherapy, educational package and drug therapies CG: routine pregnancy treatment	In the investigation of mental health subscales in the IGs and CGs, results demonstrated a significant intergroup difference in the 35–37 week follow-up in terms of anxiety ($p=.003$). Anxiety showed a significant decrease in the intervention group compared to the CG.	General Health Questionnaire-28 (GHQ-28) Golombok Rust Inventory of Marital State (GRIMS)	Mental health of pregnant women - anxiety

Uludağ (2020)	Turkey Date of data collection could not be stated RCT moderate	$N=60$ Age: IG: 25.66 ± 4.33 CG: 24.70 ± 4.75 Nulliparous	IG: Philosophy of HypnoBirthing CG: Routine Care	A statistically significant difference was found between the labor fear mean score in terms of group, time and group*time interaction ($p<.05$). There was a significant difference between the post-intervention, active phase and transition phase labor fear mean score of the groups in terms of the intervention performed: the fear of labor was lower in the IG compared to the CG.	Visual analog scale of determining the fear and pain of labor	Labor fear
Rajeswari (2020)	India 2015-2016 RCT moderate	$N=250$ Age: Majority were in the age group of 25–29 years (IG 60 [48%]; CG 57 [45.60%]). Primigravida	IG: Routine Care + progressive muscle relaxation CG: routine antenatal care	In the posttest, the groups exhibited significant difference for stress ($F_3=24.81, p<.001$) and overall anxiety ($F_3=19.80$ with $p<.001$). After the test, there was a significant reduction in state anxiety ($F_3=17.80, p<.001$) and trait anxiety ($F_3=18.60, p<.001$) between the intervention and control groups. There was a strong negative correlation between PMR and state anxiety ($r = -0.26, p<.001$).	State-Trait Anxiety Inventory (STAI)	State/ Trait Pregnancy anxiety
Bazrafsahn (2020)	Iran 2019 RCT strong	$N=72$ Age: IG: 28.06 ± 4.33 CG: 26.22 ± 4.43 Not stated	IG: group educational counseling sessions (integration of psychological instructions and interactive lectures) + routine care CG: routine pregnancy care	There was a significant difference in the mean anxiety score between the IG and CG before the group educational counseling sessions. After this intervention, a significant reduction in the mean anxiety scores of intervened pregnant women compared to the control was found. This decrease in mean anxiety score after the 1-month post-counseling was more pronounced than the 6 th week after the study onset ($p<.001$). Low anxiety scores in the intervention group over time were also maintained.	Pregnancy-related anxiety questionnaire Short-form PRAQ with 17 items (PRAQ-17)	Preg-nancy related anxiety
Munkhondya (2020)	Malawi 2018 quasi-experimental study moderate	$N=70$ Age: IG: $19.83 (\pm 2.90)$ CG: $20.11 (\pm 2.70)$ Primigravida	IG: companion-integrated childbirth preparation (structured childbirth education) CG: Routine care	At post-test, being in the intervention group significantly decreased childbirth fears ($\beta = -.866, t(68)=-14.27, p<.001$).	Childbirth Attitude Questionnaire (CAQ)	FOC
Price (2019)	USA Date of data collection could not be stated one-group repeated measures design moderate	$N=12$ Age (median, range): 30.5 (24–40) Nulli- and Multiparous	Mindfulness-Based Childbirth and Parenting (MBCP) – online audios	The significant pre-post intervention improvements included a decrease in prenatal pregnancy anxiety ($p=.002$), and increased interoceptive awareness skills of self-regulation ($p=.016$) The significant longitudinal improvements included interoceptive awareness skills of self-regulation ($p=.04$). The effect sizes for these significant improvements were large, ranging from 0.62 to 1.18.	Generalized Anxiety Disorder Scale (GAD-7)	General anxiety during pregnancy
Yang (2019)	China 2018 RCT strong	$N=123$ Age: IG: 31.31 (4.97) CG: 30.38 (3.91) Nulli- and multiparous	IG: online mindfulness intervention program (training acceptance for internal and external experiences) CG: routine prenatal care	In the IG, the mean scores of the PHQ-9 and GAD-7 before the intervention indicated mild symptoms of anxiety; these scores decreased significantly at the end of the intervention, indicating no symptoms ($t=6.218, p<.001$; $t=5.422, p<.001$, respectively). No changes in the PHQ-9 and GAD-7 scores were observed in women in the CG when scores before versus after intervention were compared. Postintervention scores of both PHQ-9 and GAD-7 were significantly lower in the IG than in the CG. Additionally, a larger proportion of women in the IG had no symptoms of anxiety after the IG compared with women in the CG.	Generalized Anxiety Disorder Scale (GAD-7)	General Anxiety during pregnancy

Van der Zwan (2019)	Netherlands Date of data collection could not be stated RCT weak	<i>N</i> =50 Age: 31.6 (5.9) Primi-/Nulli- and Multiparous	IG: heart rate variability (HRV)-biofeedback + Stress-Reducing Intervention (psycho-education + taught abdominal breathing and HRV biofeedback) CG: Waitlist condition	In both conditions anxiety and stress levels were reduced and well-being increased between pre- and post-test (T1–T2). In the HRV-biofeedback condition, within-group effect sizes were medium, and long-term improvements six weeks after the training (T1–T3) were similar to those at post-test for all outcome measures except depression. Statistically significant long-term improvements in the HRV-biofeedback condition were present for stress and psychological well-being. Effect sizes were larger in the HRV-biofeedback condition than in the waitlist condition on all outcome variables except anxiety. When comparing the treatment effect between pregnant and non-pregnant women (the Condition–Pregnancy interaction), a statistically significant interaction effect for anxiety appeared. Additional analyses showed that HRV-biofeedback was more beneficial regarding anxiety reduction for pregnant women than for non-pregnant women (pregnant women: $B=-4.18$, $t=-2.74$, $p=.006$; non-pregnant women: $B=2.55$, $t=1.99$, $p=.046$).	Dutch version of the Depression Anxiety Stress Scales (DASS)	Anxiety during pregnancy
Loughnan (2019)	Australia Date of data collection could not be stated RCT weak	<i>N</i> =77 Age: 31.61 (4.00) Primi - and Multiparous	IG: internet-delivered cognitive behavioral therapy CG: treatment as usual (TAU)	The group by time interactions for psychological distress ($F(2,53.93)=7.07$, $p<.01$) and anxiety ($F(2,54.67) = 6.48$, $p<.01$) were significant. Participants in the IG demonstrated large and superior reductions in distress at post-assessment compared to CG ($g(95\%CI) = 0.88(0.34,1.43)$), and moderate differences at follow-up, although these were not statistically significant ($g(95\%CI) = 0.52(-0.07,1.10)$). The between group differences for anxiety severity were small and non-significant post-assessment ($g(95\%CI)=0.40(-0.13,0.93)$). However, IG demonstrated a moderate to large effect size reduction in anxiety symptom severity at follow-up assessment compared to the CG ($g=0.76$; 95% CI:0.17,1.35).	Generalized Anxiety Disorder 7-item scale (GAD-7)	General Anxiety during pregnancy
Zemestani (2019)	Iran Date of data collection could not be stated RCT strong	<i>N</i> =38 Age: IG: 28.63 (3.02) CG: 30.54 (4.15) Not stated	IG: Mindfulness-based cognitive therapy (MBCT) intervention CG: Did not receive any intervention; after 1- month follow-up, two psychoeducational sessions were conducted	Results from the mixed method repeated measure (MMRM) indicate greater improvements in levels of anxiety in the IG than in the CG. As to BAI, results indicated a significant effect of time, $F=(43.72)$, $p<.0001$, $\eta^2 =.62$; and a significant time×group interaction, $F=(52.68)$, $p<.0001$, $\eta^2 =.67$. Post hoc comparisons showed that the IG had a significant decrease in BAI scores from baseline to post-treatment and BAI scores remained significantly lower than those of the CG at follow-up ($p<.0001$).	Beck Anxiety Inventory (BAI)	Anxiety during pregnancy
Amiri (2019)	Iran 2018 RCT strong	<i>N</i> =68 Age: IG: 26.2 (5.4) CG: 27.0 (5.6) Not stated	IG: Counseling based on distraction techniques for controlling stress, fear and pain CG: training about signs and stages of delivery and the appropriate time for a referral to the hospital	There was no statistically significant difference between the two groups before the intervention ($p=.117$). But in the 36th week of pregnancy the mean score of the fear of childbirth in the IG was less than that of the CG, but the difference was not statistically significant (AMD: 5.4; 95% CI: -2.4 to 13.0; $p=.117$). There was no statistically significant difference between the groups after intervention ($p=.170$).	W-DEQ-A	FOC
Rahmani (2019)	Iran Date of data collection could not be stated RCT moderate	<i>N</i> =108 Age (18-35): IG 1: 24.4 (4.14) IG 2: 26.52 (4.6) CG: 25.6 (4.35) Primi- and Multigravida	IG 1: Peer Education + training booklet IG 2: Discussion Groups + training booklet CG: not described	Significant difference among the 3 groups ($p=.007$) after 4 weeks of intervention. Further, the Scheffe test showed a significant difference between the peer education and control groups ($p=.04$), as well as the training and discussion groups with the peer education group ($p=.013$). The fears decreased in the IG1 and IG2 compared to the CG ($p<.007$) four weeks after education.	Widget's Maternity Fear Awareness Questionnaire	FOC

Zhang (2018)	China 2016 RCT moderate	<i>N</i> =66 Age: IG: 25.7(2.79) CG: 25.58(2.33) Primi- and multiparous	IG: Mindfulness stress reduction (MBSR) CG: treatment-as- usual	The results found a significant interaction between time and condition for anxiety ($F=19.30, p<.001,$ ($\eta^2=0.240$) Post hoc comparisons showed that the IG had a stronger decrease in STAI from baseline to post- treatment compared to the CG.	State Trait Anxiety Inventory (STAI)	State/ Trait Anxiety during preg- nancy
Hildingsson (2019)	Sweden 2016- 2017 Experimental Study strong	<i>N</i> =70 Age: <32: 29 (41.4) ≥ 32: 41 (58.6) Primi- and Multiparous	IG: Counseling through known midwives CG: Counseling through unknown midwives	No differences on level of fear in IG (mean FOBS 71.25; 20.41) versus CG (70.83; 21.52).	Fear of Birth Scale (FOBS)	FOC
Klabbers (2019)	Netherlands 2012-2015 RCT weak	<i>N</i> =134 Age: IG 1: 32.8 (SD 4.6) IG 2: 31.8 (SD 3.9) CG: 32.6 (SD 5.3) Primi-and Multigravida	IG 1: Haptotherapy (HT) IG 2: Psychoeducation via the Internet (INT) CG: Care as usual (CAU)	In the intention to treat analysis, only the IG1 showed a significant decrease of fear of childbirth, $F(2,99)=.321, p=.040$. In the as treated analysis, the IG1 showed a greater reduction in fear of childbirth than the other two groups, $F(3,83)=6.717, p<.001$.	W-DEQ	FOC
Irmak (2019)	Turkey 2016- 2017 RCT strong	<i>N</i> =120 Age: IG 1: 27.29 ± 3.97 IG 2: 27.51 ± 4.65 CG: 27.36 ± 4.19 Nulliparous	IG 1: Emotional freedom techniques (EFT) IG 2: breathing awareness (BA) CG: Standard care.	No significant difference in the scores for the W-DEQ-A between the groups ($p>.05$). However, the difference in the scores for the W-DEQ-B between the groups was significant ($p<.001$). This difference was due to the high score of the W-DEQ-B of the CG. Both IG1 and IG2 interventions enabled to reduce the level of birth fear perceived at postpartum. There was also a significant difference in the scores for the W-DEQ-B subscales related to worries about childbirth ($p<.05$).	W-DEQ Subjective Units of Distress Scale (SUDES)	FOC
Heller (2020)	Netherlands Date of data collection could not be stated RCT moderate	<i>N</i> =79 IG: 32.08 (4.61) CG: 31.94 (4.83) Nulli- and Multiparous	IG: internet-based problem solving treatment (PST) CG: Care as usual	In the IG, affective symptoms decreased more than in the CG, but between-group effect sizes were small to medium (Cohen's d at T3=0.45, 0.21, and 0.23 for the 3 questionnaires, respectively) and statistically not significant.	Hospital Anxiety and Depression Scale- Anxiety subscale (HADS-A)	Anxiety during pregnancy
Mohammadi (2019)	Iran 2018 RCT moderate	<i>N</i> =60 Age: IG:28.18±3.38 CG: 28.63±3.14 Not stated	IG: Intervention group attended Benson's relaxation technique (BRT) and brief psychoeducational intervention (BPI) educational sessions CG: Received no intervention	Significant statistical difference in the IG before and after intervention ($p<.001$). In the IG, the mean stress and anxiety scores, and total score were decreased significantly ($p<.001$). The CG did not show any significant statistical differences ($p>$.05). There was a significant difference between the mean scores of IG and CG ($p<.001$). In the IG the mean anxiety score significantly decreased ($p<.001$).	Depression Anxiety Stress Scale-21	Anxiety during pregnancy
Ekrami (2019)	Iran 2017 RCT strong	<i>N</i> =80 Age: IG: 28.5 (7.4) CG: 30.7 (5.4) Not stated	IG: sessions of individual counseling + sessions of group counseling CG: received routine care	The mean (SD) state anxiety score in the IG decreased from before intervention to 4 weeks after counseling; the mean (SD) state anxiety score in the CG increased from before the intervention to 4 weeks after the completion of the counseling. No significant difference between the IG and CG before the intervention in terms of state anxiety score ($p=.759$). The mean state anxiety score in the IG was significantly lower than on the CG (adjusted mean difference: -7.8, CI 95% -4.5 to -11.1; $p<.001$) after intervention. The mean (SD) trait anxiety score in the IG decreased from before counseling to 4 weeks after counseling; the mean (SD) trait anxiety score in the IG was increased from before the intervention to 4 weeks after the completion of the counseling. There was no significant difference between the IG and CG before the intervention in terms of trait anxiety score ($p=.473$). The mean trait anxiety score in the IG was significantly lower than on the CG (adjusted mean difference: -8.2, CI 95% -10.9 to -5.4; $p<.001$) after intervention.	Spielberger State- Trait Anxiety Inventory (STAI)	State/ Trait Anxiety of women with un- planned preg- nancy

Sobhani (2019)	Iran 2017 RCT weak	<i>N</i> = 40 Age could not be stated. Not stated	IG: Mindfulness Based Stress Reduction (MBSR) CG: unclear	Mindfulness training had a significant effect on reducing anxiety and stress.	Depression Anxiety Stress Scale (DASS-21)	Anxiety during pregnancy
Kacperczyk-Bartnik (2019)	Poland 2016 cross-sectional survey-based study moderate	<i>N</i> =147 Age: 31.5 (±4.8) Primi- and Multiparous	IG: Antenatal classes attendance CG: No antenatal classes attendance	Women who gave birth for the first time and attended antenatal classes scored significantly lower in the DFS questionnaire ($p<.03$). No significant differences in the DFS score were observed in case of patients giving birth for the second or subsequent time. Respondents in the IG scored slightly lower in comparison to the CG ($p<.90$).	Delivery Fear Scale (DFS)	FOC
Uçar (2019)	Turkey 2012-2013 pretest-posttest experimental design weak	<i>N</i> =111 Age: 25.5 (SD 4.2) Primigravida	IG: educational program on coping with childbirth fears based on CBT CG: did not receive any intervention	The post-education W-DEQ-A score was significant higher in the CG compared to the IG ($p<.000$). No statistically significant difference was found between the anxiety levels of the IG and CG during the active phase of labor, according to the sum of SAI scores ($p=.533$).	State Anxiety Inventory (SAI) W-DEQ-A	FOC State anxiety during pregnancy
Narita (2018)	Japan Date of data collection could not be stated Experimental Study weak	<i>N</i> =97 Age: IG: 32.4 (± 3.8) CG: 32.7 (± 5.0) Primi- and Multigravida	IG: heart rate variability (HRV) biofeedback Intervention (Stress Eraser) CG: women did not agree to practice the method	The W-DEQ scores reduced significantly in women who performed HRV biofeedback ($n=18$, $p<.001$), but there was no change in those who did not perform the method ($n=20$).	W-DEQ-A	FOC
Boryri (2018)	Iran 2017 Quasi Experimental Study moderate	<i>N</i> =180 Age: 24.54 ± 4.40 Primiparous	IG 1: muscle relaxation IG 2: guided imagery CG: Routine care	The scores of delivery fear before the intervention significantly differed in the three groups ($p=.01$). A significant difference was found between IG1 and IG2 ($p=.01$), while the other groups represented no difference. However, the mean score of the fear of delivery was significant in the three groups after the intervention ($p=.0001$). The post-hoc test further indicated a statistically significant difference in the mean scores of childbirth fear between the IG1 and IG2 ($p=.0001$), IG1 and CG ($p=.0001$), as well as IG2 and CG ($p=.0001$). In the IG1 and IG2 fear of delivery was reduced significantly.	Brislin's questionnaire	FOC
Warriner (2018)	UK 2014-2015 initial pilot study weak	<i>N</i> =155 (86 women, 69 men) Age (mean): 35 years Not stated	IG: 'MBCP-4-NHS' - Brief four week course (developed from the nine week Mindfulness Based Childbirth and Parenting (MBCP) intervention)	Change in mood pre-to post course showed that all scores improved and were statistically significant for prospective mothers, except for positive pregnancy experience intensity. Anxiety score has reduced to the 'mild' cut-off.	Generalized Anxiety Disorder Scale (GAD-7) Oxford Worries about Labor Scale (OWLS) Pregnancy Experience Scale (PES) Brief Tilbury Pregnancy Distress Scale (TPDS)	General anxiety during pregnancy Worries about labor Preg-nancy distress
Akbarian (2018)	Iran 2016 RCT weak	<i>N</i> =120 Age was not stated. Primiparous	IG: couples (mental health training course; with the partner present), pregnant women (mental health training course without the partner present) CG: routine care	In the pregnant women group and couples group, the average anxiety score of pregnant women after the intervention was significantly lower than before the intervention ($p<.001$). A significant difference was shown among the three groups after the intervention. After the intervention, the mean anxiety score of the pregnant women group was significantly lower than that of the CG ($p=.002$) and this score was significantly lower in the couples group than that in the pregnant women group ($p=.045$).	Depression, Anxiety, and Stress Scales (DASS-42)	Anxiety during pregnancy

Krusche (2018)	UK Date of data collection could not be stated RCT moderate	$N=185$ Age (mean): 32.7 Primi- and multiparous	IG: online mindfulness course ('Be Mindful Online') - immediate CG: waiting to take the mindfulness course after the baby was born	A pairwise comparisons showed a decrease in anxiety for immediate, $[F(1,69) = 18.42, p < .001, \eta^2 = .21]$ (mean difference -3.88) and waitlist participants, $[F(1,69) = 14.27, p < .001, \eta^2 = .17]$ (mean difference -2.23). There was a trend for immediate participants to have lower anxiety at T1 compared to waitlist controls, $[F(1,69) = 3.15, p = .08, \eta^2 = .04]$.	The General Anxiety Disorder-7 (GAD-7)	General anxiety during pregnancy
Rondung (2018)	Sweden 2014-2015 RCT moderate	$N=258$ Age: <25: 37 (14.3) 25-35: 186 (72.1) >35: 35 (13.6) Primi- and multiparous	IG: Guided internet-based on cognitive behavioral therapy (ICBT) CG: Standard care group	The reduction in FOB over time was significantly larger in the guided IG group than in the CG group. However, the predicted level of FOB at the estimated due date did not differ significantly ($t_{1,240,996} = -0.24, p = .81$). Hence, when comparing the intervention groups, no difference was observed in FOB in late pregnancy.	Fear of Birth Scale (FOBS)	FOC
Hajmohamadi (2018)	Iran 2014 RCT moderate	$N=114$ Age was not stated Not stated	IG: Psycho-education CG: Not stated	The mean score of depression and anxiety decreased significantly after the intervention in comparison to that before the intervention and that of CG ($p < .001$).	Researcher made questionnaire based on the Predisposing, Reinforcing and Enabling Constructs in Educational Diagnosis and Evaluation	Anxiety during pregnancy
Airo (Toivanen) (2018)	Finland 2007-2010 Randomized trial weak	$N = 460$ Age: IG: 29.8 (± 4.4) CG: 28.3 ± 5.0 Primiparous	IG: group intervention Nyytti® (with psychoeducation elements, the lifespan model of motivation, practices to support mentalisation and mind-body connection).	FOC decreased statistically significant during the intervention from the baseline (mean=7.60, SD=1.72) to the last session before childbirth (sixth session; mean=4.56, SD=2.42, Wald=230.43, df=6, $p < .001$).	W-DEQ-A Visual Analog Scale (VAS) to measure subjective FOC	FOC
Duncan (2017)	USA Date of data collection could not be stated RCT moderate	$N=30$ Age could not be stated Nulliparous	IG: Short, time-intensive course. Mind in Labor (MIL): Working with Pain in Childbirth, based on Mindfulness-Based Childbirth and Parenting (MBCP) education CG: standard childbirth preparation course with no mind-body focus	Pain catastrophizing dropped by 3.6 points in the IG group and was essentially unchanged in the CG. The time*group interaction was not significant ($t = -1.06, p = .30$; estimated treatment effect = -3.26 points, 80% CI [-7.3, 0.8]). When the missing data was imputed, the result did not change ($t = -.71, p = .48$).	Pain Catastrophizing Scale (PCS)	Pain catastrophizing
Andaroon (2017)	Iran 2015-2016 RCT moderate	$N=93$ Age could not be stated Primiparous	IG: face to face individual counseling CG: Usual services	The present study showed that an individual counseling program provided by a midwife based on a counseling consultant by a midwife based on BELIFE counseling is effective in reducing fear of childbirth in a way that the level of fear of childbirth in primiparous women in the weeks 36-34 of pregnancy in the IG was significantly lower than the CG.	W-DEQ	FOC
Seyed Kaboli (2017)	Iran 2016 RCT strong	$N=62$ Age: 19 (30.64) 29-25: 29 (46.77) 35-30: 14 (22.58) Primiparous	IG: counseling for 6 sessions of 90 minutes + routine prenatal care CG: routine prenatal care + instructional package for dealing with pregnancy stresses	The PWSQ score did not differ significantly between the 2 groups before the intervention ($p > .05$). After the intervention, the mean subscale scores were lower in the IG than in the CG and showed a statistically significant post-intervention difference between the groups ($p = .01$). These scores suggest the effectiveness of the intervention in reducing pregnancy-specific stress.	Pregnancy Worries and Stress Questionnaire (PWSQ)	Preg-nancy worries and stress

Mary (2017)	India Date of data collection could not be stated pre- test, post-test quantitative research design weak	<i>N=50</i> Age: 64 % of participants in CG and IG were between 24-29 years Primi antenatal	IG: performed selected mind body interventions (Active visualisation with Birth Affirmations, yogic breathing and relaxation) for 4 weeks CG: routine standard hospital care	Statistical findings proved that there was a significant difference in anxiety level among antenatal women who were subjected to mind body intervention than those who were not.	W-DEQ	FOC
Legrand (2017)	France Date of data collection could not be stated single-subject A (baseline) – B (hypnotherapy treatment) – A' (return-to-baseline) research design weak	<i>N=1</i> Age: 23 years Primigravida	Hypnotherapy treatment	A statistically significant declining trend in anxiety scores was observed during the hypnosis phase, and anxiety re-increased in the return-to-baseline phase ($p<.05$).	State Anxiety Inventory (SAI)	State anxiety during pregnancy
Waisblat (2017)	France Date of data collection could not be stated longitudinal repeated measures quasi-experimental design moderate	<i>N=155</i> Age: Group S: 44.3 (13.3) Group H: 46.3 (7.1) Not stated	Group S: standard hypnotic communication Group H: hypnotic communication	The mean fear ratings in the Group H participants were significantly lower than that of the Group S participants ($p=.001$).	Rating fear of the epidural puncture using the numerical rating scale (NRS) with 0 = No pain (fear) and 10 = Worst imaginable pain (fear).	Fear of women under-going labor
Toosi (2017)	Iran Date of data collection could not be stated semi-experimental clinical trial moderate	<i>N=80</i> Age: IG: 29.0 ± 2.4 CG: 28.7 ± 2.7 Primiparous	IG: Relaxation Training (Benson's relaxation technique) CG: Routine care	No significant difference between the two groups regarding the anxiety score before the intervention ($p=.903$). A statistically significant difference was observed regarding the anxiety score after the intervention ($p<.001$). The anxiety score had significantly decreased in the IG ($p<.001$), but had significantly increased in the CG ($p=.033$). Thus, relaxation training was effective in reduction of anxiety score after the intervention.	Spielberger's state-trait anxiety scale	State/Trait anxiety during pregnancy
Sanaati (2017)	Iran 2015 RCT strong	<i>N=189</i> Age: IG1: 28.2 (5.1) IG2: 27.5 (4.9) CG: 27.7 (4.9) Not stated	Lifestyle based education: included issues related to sleep, hygiene, nutrition, physical activity and exercise, self-concept and sexuality IG 1: both women and their husbands received the lifestyle-based education. IG2: only women received the lifestyle-based education. CG: Routine care	No significant difference on state or trait anxiety was observed between the groups before the intervention ($p=.257$; $p=.137$) The mean score of state anxiety 8 weeks after intervention showed a statistically significant difference among the groups ($p<.001$). The mean state anxiety scores in the IG1 and IG2 were significantly reduced compared to the CG. The mean state anxiety score was also significantly reduced in the IG1 compared to the IG2. The mean post-intervention score of trait anxiety showed a statistically significant difference among the groups ($p<.001$). Compared to the CG, the mean trait anxiety score was significantly reduced in the IG1 and IG2; however, no significant difference was observed between the two intervention groups.	Spielberger State-Trait Anxiety Inventory (STAI)	State/Trait Anxiety during pregnancy
Kordi (2017)	Iran 2015-2016 RCT strong	<i>N=122</i> Age: IG: 23.2 ± 3.6 CG: 24.2 ± 4.4 Primigravida	IG: psycho-educational program for three weeks CG: Routine prenatal care	No significant differences between the groups with respect to the mean pre-intervention FOC scores. A significant difference was observed between the IG and CG in terms of the mean post-intervention FOC scores ($p=.007$). The FOC score significantly diminished in the intervention group in the post-intervention phase ($p=.001$).	W-DEQ	FOC

Beattie (2017)	Australia 2014 Pilot randomized trial moderate	<i>N</i> =48 Age: IG: 28.9 (5.7) CG: 28.5 (6.4) Nulli- and Multiparous	IG: mindfulness-based support program (MiPP) CG: pregnancy support program (PSP)	No statistically significant differences between the IG and the CG were shown on perceived stress across the three time periods (<i>p</i> =.82).	Perceived Stress Scale (PSS-10) (PSS)	Perceived stress during pregnancy
Haapio (2017)	Finland Date of data collection has not been stated RCT weak	<i>N</i> =659 Age (%): 18–22: IG: 3.0%; CG: 4.0% 23–29: IG: 48.0%; CG: 47.0% 30–35: IG 44.0%; CG 45.0% 36–40: IG: 5.0 %; CG: 4.0% Primiparous	IG: extended childbirth education (defined as a midwife-led intervention with low medicalization) CG: regular childbirth education	The mothers in the IG had less childbirth-related fear than those in the CG [odds ratio (OR) 0.58, 95% confidence level (CL) 0.38–0.88].	'Feelings of Fear and Security Associated with Pregnancy and Childbirth' Questionnaire	FOC
Parsa (2016)	Iran 2015 Quasi experimental study strong	<i>N</i> =110 Age (IG/CG): 18-22: 16.4%/5.5% 23-27: 47.3%/49.1% 28-32: 29.1%/36.4% 33-37: 7.3%/ 9.1% Nulliparous	IG: counseling sessions based on the GATHER approach CG: not described	Trait anxiety levels of pregnant women significantly changed (were lowered) as a result of intervention (<i>p</i> <.001). However, no significant difference was found in trait anxiety levels of pregnant women in the CG before and after the intervention. State anxiety levels of pregnant women significantly changed (were lowered) as a result of intervention (<i>p</i> <.001). However, no significant difference was found in state anxiety levels of pregnant women in the CG before and after the intervention.	Spielberger's State-Trait Anxiety Inventory	State/Trait Anxiety during pregnancy
Aslami (2016)	Iran 2015 RCT moderate	<i>N</i> =75 Age: IG 1: 29.4±3.8 IG 2: 27±3.2 CG: 28.6±4.3 Not stated	IG1: treatment of mindfulness based on Islamic spiritual schemes IG2: cognitive behavioral therapy group CG: no course	The significant levels of all tests reveal that between the anxiety of pregnant women in the IGs and CG, at least in one of the dependent variables in the <i>p</i> <.001 level, there was a significant difference. This finding shows that in the aforesaid related variables statistically significant differences are seen between IG1 and IG2 and CG. These findings revealed that both IG1 and IG2 in comparison to the CG led to a decrease in anxiety in pregnant women. The difference between the average IG1 and IG2 in anxiety was significant in the level of <i>p</i> <.001. Therefore, the mindfulness treatment method in comparison with group cognitive behavioral therapy was more effective on the reduction of anxiety of pregnant women.	Beck anxiety-depression questionnaire	Anxiety during pregnancy
Khojasteh (2016)	Iran 2016 RCT moderate	<i>N</i> =75 Age: IG1: 22.76 ± 3.85 IG2: 23.76 ± 3.74 CG: 23.92± 4.41 Nulliparous	IG1: Massage IG2: Guided Imagery CG: Routine Care	No significant difference before intervention between groups (<i>p</i> =.063). The mean score of anxiety in all three groups had statistically significant differences after the intervention (<i>p</i> <.001). The post-hoc test showed that the IG1 (<i>p</i> =.000) and IG2 (<i>p</i> =.000) had significant lower anxiety scores compared to the CG, while no significant differences were found between IG1 and IG2 (<i>p</i> =.928).	Pregnancy-related Anxiety Questionnaire - revised	Pregnancy related anxiety
Sheikh-Azadi (2016)	Iran Date of data collection could not be stated RCT moderate	<i>N</i> =60 Age: IG: 24 (4.388) CG: 25 (4.387) Not stated	IG: Routine pregnancy care + group discussion courses CG: Routine pregnancy care	Mean anxiety score before the intervention was not significantly different between the IG and CG (<i>p</i> =.674). The results showed, that the mean anxiety score of maternal state anxiety was significantly different between the two groups after the intervention (<i>p</i> =.001). It was significantly lower in the IG compared to the CG.	Spielberger Anxiety Inventory	State anxiety during pregnancy

Salehi (2016)	Iran 2015 quasi experimental trial strong	<i>N</i> =91 Age: 26.04±4.68 Nulliparous	IG1: group cognitive behavioral therapy (GCBT) IG2: interactive lectures group (IL) CG: standard prenatal care	There was a significant difference in the level of state and trait anxiety in both the IG1 and IG2 groups before and after the intervention ($p<.001$). However, there were no differences in state anxiety ($p=.330$) or trait anxiety ($p=.147$) in the CG between baseline and 4 weeks later. The results showed significant differences between the 3 groups in state anxiety ($p=.011$) and trait anxiety ($p=.016$). No significant difference was found between IG1 and IG2 for state anxiety ($p=.079$) or trait anxiety $p=.069$. GCBT and IL significantly reduced anxiety in pregnant women.	Spielberger's State-Trait Anxiety Inventory	State/ Trait Anxiety during pregnancy
Beevi (2016)	Malaysia Date of data collection could not be stated pre-test/post-test quasi- experimental design moderate	<i>N</i> =56 Age: IG: M = 28.23 SD = 3.12 CG: (M = 29.28 SD = 2.65) Nulli- and Multiparous	IG: Hypnosis intervention CG: Traditional antenatal care	There was a statistically significant interaction between the group and time for anxiety symptoms, $F(3,126)=7.933, p<.037$, partial $\eta^2=.16$. Results for the simple main effect for group indicated that there was a statistically significant difference in anxiety symptoms at time point 3, $F(1,44)=10.764, p=.002$, partial $\eta^2=.20$, but not at baseline, time point 1 and time point 2. There was a statistically significant effect of time on anxiety symptoms for the IG, $F(2.138,58.457)=12.352, p=.0005$, partial $\eta^2=.38$ and the effect of time on anxiety symptoms for the CG was not significant, $F(3,66) = 0.756, p=.523$, partial $\eta^2 =.03$. Following the significant effect of time for the IG, a pairwise comparison was performed and results indicated that anxiety symptoms were statistically significantly reduced between time point 1 and baseline ($M=2.48, SE=0.80, p=.035$), between time point 2 and baseline ($M=3.91, SE=1.18, p=.020$), between time point 3 and baseline ($M=6.10, SE=1.27, p=.001$), between time point 3 and time point 1 ($M=3.62, SE=1.13, p=.026$), but not statistically significant between time point 1 and time point 2 ($M=1.43, SE=0.89, p=.734$) and between time point 2 and time point 3 ($M=2.19, SE=0.82, p=.085$)	Depression Anxiety Stress Scale—21 (DASS-21)	Anxiety during pregnancy
Fontein-Kuipers (2016)	Netherlands 2013-2015 RCT strong	<i>N</i> =433 Age: IG: 30.11 (±4.09) CG: 29.98 (±3.71) Nulli- and multiparous	IG: Wazzup Mama?! focused 1. on the signs and symptoms of maternal distress and identification of the origin of the state of mood 2. identifying stressors 3. measurement of maternal distress. CG: antenatal care as usual	In the CG, the mean STAI scores significantly increased from baseline (T1) to post-intervention (T2) ($p<.001, p<.001, p<.001$). Mean PRAQ scores increased but did not reach statistical significance ($p=0.12$). The proportion of STAI and PRAQ scores above cut- off level significantly increased from baseline (T1) to post-intervention (T2) ($p<.001, p=.045, p=.03$). In the IG, the mean STAI and PRAQ scores were significantly lower at T2 compared to T1 ($p=.001, p<.001, p<.001$). The proportion of PRAQ scores above cut-off level were significantly lower at T2 compared to T1 ($p=.002, p=.009$). The STAI scores above cut-off level decreased, but this did not reach statistical significance ($p=.4, p=.4$).	State-Trait Anxiety Inventory (STAI) and Pregnancy-Related Anxiety Questionnaire (PRAQ)	State/ Trait Anxiety in pregnancy
Yazdanimehr (2016)	Iran Date of data collection has not been stated RCT strong	<i>N</i> =80 Age: IG: 26 (5.82) CG: 26.73 (4.54) Not stated	IG: Mindfulness-integrated cognitive behavior therapy CG: Routine prenatal care services	The differences between the study groups regarding the pretest mean scores of anxiety were not statistically significant ($p<.05$). The results showed that at T2 and T3, the mean scores of anxiety in the IG were significantly lower than the CG ($p<.001$).	Beck Anxiety Inventory	Anxiety during pregnancy
Nieminen (2016)	Sweden 2012- 2013 feasibility study weak	<i>N</i> =28 Age: 30.5 (24-39) Nulliparous	IG: Internet-delivered therapist-supported self-help program based on cognitive behavioral therapy (ICBT)	Statistically significant ($p<.0005$) decrease of FOC. The W-DEQ sum score decreased pre- to post-therapy, with a large effect size (Cohen's $d=0.95$).	W-DEQ	FOC

Karabulut (2016)	Turkey date of data collection has not been stated quasi-experimental & prospective study strong	N=192 Age: IG: 28.87 ± 4.54 CG: 25.73 ± 5.35 Primiparous	IG: Antenatal educational program (health in pregnancy, birth and breathing exercises, breastfeeding, baby care, post-partum period and family planning) CG: routine pregnancy care and information	The IG's pre-education and IG's first measurement levels of FOC showed significant differences ($p<.005$). The IG's post-education and CG's second measurement levels of FOC also showed significant differences ($p<.005$). According to this finding, antenatal education was effective in reducing the FOC among primipara.	W-DEQ-A	FOC
Karamoozian (2015)	Iran Date of data collection has not been stated RCT moderate	N=30 Age is not reported Primiparous	IG: cognitive-behavioral stress management (CBSM) CG: prenatal care	There is a significant difference in the adjusted average scores of total anxiety between the IG and CG. The effect of pretest was significant with $\eta^2_p=0.57, p<.01$, and $f=34.83$. As a result, it can be said that CBSM significantly reduced the total anxiety in the IG.	Pregnancy-Related Anxiety Questionnaire	Preg-nancy related anxiety
Rouhe (2015)	Finland 2007-2009 RCT Weak	N=371 Age not stated Not stated	IG: group psycho-education with relaxation exercises CG: conventional care	There was a significant difference between the groups in mean W-DEQ-B sum scores ($F=1.1, df=199, p=.016$ Cohen's $d=0.35$, small effect size), indicating a more fearful childbirth experience in the CG. Childbirth experience was less fearful in the IG compared to the CG across all modes of delivery, although none of the differences reached significance, potentially because of small sample sizes.	W-DEQ A+B	FOC
İsbir (2015)	Turkey 2014 RCT strong	N=72 Age: IG: 24.9 (5.9) CG: 25.0 (4.7) Primi- and multiparous	IG: Supportive Care during labor by midwives (physical, emotional, instructional, informational, advocacy support) CG: routine hospital care	The IG reported less fear of delivery during the active and transient phases of labor than the CG ($p<.05$).	W-DEQ "Delivery fear scale"	Fear of delivery

Table 1: Summary of the included studies (Description of Studies).

		Selection Bias	Study Design	Confounders	Blinding	Data collection method	Withdrawals and dropouts	Global rating of paper
Aksoy (2021)	Derya	moderate	strong	moderate	moderate	strong	weak	moderate
Montazeri (2020)		moderate	strong	strong	weak	strong	strong	moderate
Waters (2020)		moderate	weak	weak	moderate	weak	weak	weak
Zarenejad (2020)		moderate	strong	strong	moderate	strong	strong	strong
Firouzan (2020)		strong	strong	strong	moderate	strong	strong	strong
Kang (2020)		moderate	strong	weak	moderate	strong	weak	weak
Goetz (2020)		moderate	moderate	weak	moderate	strong	weak	weak
Alipour (2020)		moderate	strong	strong	strong	strong	weak	moderate
Sridhar (2020)		moderate	strong	strong	strong	strong	moderate	strong
Esfandiari (2020)		moderate	strong	weak	strong	strong	moderate	moderate
Mirtabar (2020)		moderate	strong	strong	strong	strong	strong	strong
Abbasi (2020)		strong	strong	strong	weak	strong	strong	moderate
Larsson (in press)		weak	strong	strong	weak	weak	weak	weak
Swift (in press)		moderate	strong	strong	strong	moderate	moderate	strong
Shahsavvan (2020)		strong	moderate	strong	moderate	strong	strong	strong
Boz (2020)		moderate	strong	strong	moderate	strong	weak	moderate
Abdollahi (2020)		weak	strong	strong	moderate	strong	strong	moderate
Hamilton (2020)		weak	strong	weak	weak	strong	weak	weak
Noorbala (2019)		moderate	strong	strong	moderate	strong	weak	moderate
Uludağ (2020)		moderate	strong	strong	moderate	weak	strong	moderate
Rajeswari (2020)		moderate	strong	weak	moderate	strong	strong	moderate
Bazrafshn (2020)		strong	strong	strong	moderate	strong	strong	strong

		Selection Bias	Study Design	Confounders	Blinding	Data collection method	Withdrawals and dropouts	Global rating of paper
	Munkhondya (2020)	moderate	strong	strong	moderate	strong	strong	strong
	Price (2019)	moderate	moderate	weak	moderate	strong	strong	moderate
	Yang (2019)	moderate	strong	strong	moderate	strong	strong	strong
	van der Zwan (2019)	moderate	strong	strong	weak	strong	weak	weak
	Loughnan (2019)	moderate	strong	weak	moderate	strong	weak	weak
	Zemestani (2019)	strong	strong	strong	moderate	strong	strong	strong
	Amiri (2019)	moderate	strong	strong	moderate	strong	strong	strong
	Rahmani (2019)	strong	strong	strong	weak	strong	strong	moderate
	Zhang (2019)	moderate	strong	strong	moderate	strong	weak	moderate
	Hildingsson (2019)	weak	strong	strong	moderate	strong	strong	strong
	Klabbers (2019)	strong	strong	strong	weak	strong	weak	weak
	Irmak Vural (2019)	moderate	strong	strong	moderate	strong	strong	strong
	Heller (2020)	moderate	strong	weak	moderate	strong	moderate	moderate
	Mohammadi (2019)	strong	strong	strong	weak	strong	strong	moderate
	Ekrami (2019)	moderate	strong	strong	moderate	strong	strong	strong
	Sobhani (2019)	moderate	strong	weak	moderate	strong	weak	weak
	Kacperczyk-Bartnik (2019)	moderate	weak	strong	moderate	strong	strong	moderate
	Uçar & Golbasi (2019)	weak	moderate	strong	moderate	strong	weak	weak
	Narita (2018)	moderate	moderate	strong	weak	strong	weak	weak
	Boryri (2018)	moderate	moderate	strong	moderate	strong	weak	moderate
	Warriner (2018)	moderate	weak	weak	moderate	strong	weak	weak
	Akbarian (2018)	moderate	strong	weak	moderate	strong	weak	weak
	Krusche (2018)	strong	strong	strong	moderate	strong	weak	moderate
	Rondung (2018)	moderate	strong	strong	strong	weak	strong	moderate
	Hajmohamadi (2018)	strong	strong	strong	moderate	weak	strong	moderate
	Airo (Toivanen) (2018)	weak	strong	weak	moderate	weak	strong	weak
	Duncan (2017)	moderate	strong	weak	moderate	strong	strong	moderate
	Andaroon (2017)	strong	strong	weak	moderate	strong	strong	moderate
	Seyed (2017)	moderate	strong	strong	moderate	strong	strong	strong
	Mary (2017)	moderate	moderate	weak	moderate	strong	weak	weak
	Legrand (2017)	weak	weak	-	-	strong	moderate	weak
	Waisblat (2017)	strong	moderate	strong	moderate	weak	strong	moderate
	Toosi (2017)	moderate	strong	strong	strong	moderate	strong	moderate
	Sanaati (2017)	strong	strong	strong	moderate	strong	strong	strong
	Kordi (2017)	moderate	strong	strong	moderate	strong	strong	strong
	Beattie (2017)	moderate	strong	strong	strong	strong	weak	moderate
	Haapio (2017)	weak	strong	strong	moderate	weak	moderate	weak
	Parsa (2016)	moderate	moderate	strong	moderate	strong	strong	strong
	Aslami (2016)	moderate	moderate	strong	moderate	strong	weak	moderate
	Khojasteh (2016)	moderate	strong	strong	moderate	strong	weak	moderate
	Sheikh-Azadi (2016)	moderate	strong	strong	moderate	strong	weak	moderate
	Salehi (2016)	strong	strong	strong	moderate	strong	strong	strong
	Beevi (2016)	moderate	moderate	strong	moderate	strong	weak	moderate
	Fontein-Kuipers (2016)	moderate	strong	strong	moderate	strong	moderate	strong
	Yazdanimehr (2016)	moderate	strong	strong	moderate	strong	moderate	strong
	Nieminen (2016)	moderate	moderate	weak	weak	strong	weak	weak
	Karabulut (2016)	moderate	moderate	strong	moderate	strong	moderate	strong
	Karamoozian (2015)	moderate	strong	strong	moderate	strong	weak	moderate
	Rouhe (2015)	weak	strong	strong	moderate	strong	weak	weak
	İsbir (2015)	moderate	strong	strong	moderate	strong	strong	strong

Table 2: Ratings of included studies based on the “Quality Assessment Tool for Quantitative Studies” of the “Effective Public Health Practice Project (EPHPP)”

128,130]. Online “mindfulness based interventions” seem to be effective online/digital [96,126,128,130]. For the offline „mindfulness-based interventions“ results are inconsistent: Several studies find a positive effect, while Beattie et al. did not show a positive effect of mindfulness-based interventions on perceived stress and Duncan et al. did not show a positive result on pain catastrophizing [103,112,121,124,125,129,136,139]. This result does not seem to be affected by the weak ratings of the studies from Goetz et al., Sobhani et al. and Warriner et al. [96,121,129].

There are also mixed results concerning the effectiveness of internet based cognitive behavioral therapy. While Larsson et al. and Loughnan et al. did not find a between group effect for internet based cognitive behavioral therapy, Nieminen et al., Rondung et al. and Shahsavan et al. showed significant effects in favour of internet based cognitive behavioral therapy [83,87,89,91,127]. This result has to be interpreted carefully as the studies from Larsson et al., Loughnan et al. and Nieminen et al., were rated as weak [83,87,127].

Fontein-Kuipers focused on identifying (potential) stress factors, problems or difficult situations in the past or present that may contribute to the development of maternal distress plus gave personal feedback regarding questionnaire results in a web-based tailored program [95]. In the intervention group, the mean state anxiety and pregnancy anxiety scores were significantly lower at T2 compared to T1. The proportion of pregnancy anxiety scores above cut-off level were significantly lower at T2 compared to T1 and the state trait anxiety scores above cut-off level decreased, but this did not reach statistical significance.

Categories of interventions: Summarized within this review are *educational interventions*: psychoeducation, and more general education [74,76,80-82,85,92,93,108,109,117,131]. Bazrafshan et al, Boz et al., Hajmohamadi et al., Kordi et al. and Uçar and Golbasi found a positive effect for psychoeducation, while the study of Klabbbers did not validate this result [74,81,82,92,109,117]. Taking the ratings of Uçar and Klabbbers as weak ratings into account, the results could be interpreted as positive effects for psychoeducational interventions [81,92]. Abbasi et al., Haapio et al., Karabulut et al., Munkondhya et al. and Noorbala et al. showed a positive effect of general education, but the study of Aksoy Derya is contrary to this result [76,80,85,93,108,131]. The weak rating of the study from Haapio et al. does not seem to affect this result. One study examined the effect of partly psychoeducational, but mostly physiological education and only found a small effect of education on fear of childbirth, while Rahmani et al. showed that peer education is effective in decreasing FOC in pregnant women [47,76,88].

With regards to relaxation trainings two studies show a positive effect of *relaxation techniques* like progressive muscle relaxation and relaxation training [99,102].

The included studies show mixed results regarding *mindfulness-based interventions* and their positive effect on anxiety/fear and stress. While several authors found positive influences, two studies did not show positive effects

of mindfulness-based interventions on perceived stress or on pain catastrophizing [96,103,112,121,125,126,128-130,136,139]. This result does not seem to be affected by the weak ratings of the studies from Goetz, Sobhani and Warriner [96,121,129].

Four studies included in this review examined the effect of *hypnotherapeutic interventions*. Beevi et al. stated that an hypnotherapeutic intervention has a positive effect on reducing anxiety during pregnancy [116]. Legrand et al. also found a positive effect on decreasing state anxiety and also showed a re-increase in the return-to-baseline phase, but this study has to be interpreted carefully, as only one person was examined and the rating of the study was weak [105]. Waisblat et al. examined the effect of hypnotic communication on fear of women undergoing labor and found that hypnotic communication (communication that focusses on the awareness of the patient towards sensations and images that support relaxation and comfort) was more effective than standard communication [135]. In addition fear of labor was significantly lower in a “philosophy of hypnobirthing” group compared to the control group (received routine care) [134].

Boryri et al. and Khojasteh et al. studied the effect of *guided imagery* on FOC and pregnancy related anxiety and found a significant decrease of fear of delivery though guided imagery [73,111].

Narita et al. studied the effect of a heart rate variability (HRV) *biofeedback intervention* on fear of childbirth and found that FOC was significantly reduced in women who performed HRV biofeedback [86]. Contrary to this result, van der Zwan et al., who studied a heart rate variability (HRV)-biofeedback intervention combined with a stress-reducing intervention, did not find significant long-term improvements in the HRV-biofeedback condition [123]. But the results on both of those biofeedback intervention studies have to be interpreted carefully due to weak ratings.

Seven studies examined the effect of *counselling* on anxieties and fears related to pregnancy and childbirth. Seyed Kaboli et al. showed an effect of counselling on reducing pregnancy-specific stress. Another study, that studied face to face individual counselling conducted by a midwife was effective in reducing fear of childbirth [137,72]. Ekrami et al. examined individual and group counselling [94]. The authors found that the mean state and trait anxiety score of the counselling groups were significantly reduced compared to the control group without counselling [94]. In addition Hildingsson et al. found that it does not make a difference if the counselling is done by a known or unknown midwife. Counselling based on distraction techniques did not show a significant difference compared to a control group intervention (training about signs and stages of delivery and the appropriate time for a referral to the hospital) [71,77]. Parsa et al. examined counselling sessions based on the GATHER approach and showed that trait and state anxiety levels were lowered due to the intervention [98]. Esfandiari et al. showed that group supportive counselling scores of state-anxiety were reduced more remarkably than in the CG with a large effect size [104]. Firouzan examined the

difference between face to face counselling and telephone counselling sessions and found that counselling based on the BELIEF protocol was effective in decreasing childbirth fear [75].

This systematic review also included studies about different *therapy tools*. Montazeri et al. showed a significant and reducing effect of *writing therapy* sessions on anxiety during pregnancy [120]. An *acceptance and commitment therapy (ACT)* intervention studied by Waters et al. showed a positive effect on global distress, but must be interpreted carefully due to weak ratings [132]. Alipour et al. examined the effect of a *communication skills training* package combined with a couple based intervention as significantly effective in the reduction of anxiety during pregnancy [114]. A *cognitive analytic therapy* intervention examined by Hamilton et al. did not show any difference in trait/state anxiety between the randomized groups, but this result has to be interpreted carefully due to a weak rating [97]. Mirtabar et al. examined the effect of *individual structured psychotherapy* on state anxiety in preterm labor and showed a significant improvement in the state-anxiety scores compared with the control group (received inpatient medical care for preterm labor) [106]. Aslami et al. (2016) studied the effect of a *cognitive behavioral therapy* group on anxiety during pregnancy and revealed that the cognitive behavioral therapy group in comparison to the control group (no intervention course) led to a decrease in anxiety in pregnant women [115]. This matches the result of Salehi which also studied the effect of *group cognitive behavioral therapy (GCBT)* on state/trait anxiety during pregnancy [100]. There was a significant decrease in the level of state and trait anxiety in the GCBT group before and after the intervention. A study about a *cognitive behavioral stress management intervention* showed that this intervention significantly reduced the total anxiety [110].

Also calming *virtual reality environments* seem to be effective on reducing anxiety during pregnancy, but this effect was mainly seen by qualitative data of the study [122]. A *motivational interviewing (MI)* psychotherapy intervention showed a large and significant effect on the reduction of fear of childbirth [48]. Irmak Vural and Aslan examined the effect of *emotional freedom techniques and breathing awareness*, both interventions enabled to reduce the level of worries about childbirth [78]. An *internet based problem solving treatment* studied by Heller et al. did not show a significant effect in reducing anxiety during pregnancy [118]. *Antenatal class attendance* reduced delivery fear significantly in first time mothers, but not mothers giving birth for the second or subsequent time [79]. A *mental health training course* with and without the partner present studied by Akbarian showed that after the intervention, the mean anxiety score of the pregnant women group was significantly lower than that of the control group and this score was significantly lower in the couples group than that in the pregnant women group [113]. This result has to be interpreted carefully, due to a weak rating. Another study examined the *group intervention Nyytti®* (with psychoeducation elements, the lifespan model of motivation, practices to support mentalisation and mind-body connection) and showed a significant decrease of

fear of childbirth, but has to be interpreted carefully due to weak ratings [70]. İsbir and Serçekus studied the effect of *supportive care during labor* by a midwife (physical, emotional, instructional, informational, advocacy support) and found that women supported by the midwives showed significantly less fear on delivery [138]. A *lifestyle based education* (included issues related to sleep, hygiene, nutrition, physical activity and exercise, self-concept and sexuality) found a significant and positive effect on reducing state and trait anxiety during pregnancy [101].

Kang et al. examined the effect of different psychological interventions (like psychological support, education, relaxation training, family support, music listening) on anxiety of pregnant women undergoing interventional prenatal diagnosis [133]. The authors found that anxiety scores were significantly lower in the psychological intervention than in the control group, but the results have to be interpreted carefully due to weak ratings [133]. A *mind body intervention* on the effect of FOC showed a significantly reducing effect on the level of anxiety and anxiety symptoms among antenatal women who were assigned to mind body interventions than those who were not, but this effect has to be interpreted carefully due to a weak rating [84]. Another study focused on the effect of *group discussion* in combination with routine care with a significant decrease of state anxiety during pregnancy [107].

Discussion

Summary of main findings

This systematic review found a positive effect of *psychoeducation, relaxation techniques, guided imagery, counselling* (face to face individual counselling from a known or unknown midwife; group (supportive) counselling; counselling based on the GATHER approach or BELIEF protocol) and *different hypnotherapeutic techniques* on different fears and anxieties during pregnancy and childbirth, in the sense that those interventions have a reducing effect on fears and anxieties in the time of pregnancy and childbirth. Counselling based on distraction techniques did not show a significant difference compared to a control condition.

For *mindfulness-based interventions*, mixed results are found, regarding the positive effect of those interventions. Specially an effect on perceived stress and pain catastrophizing could not be shown.

This systematic review also included studies on *different therapeutic schools and therapy tools*, showing a significant and reductional effect of *writing therapy sessions, cognitive behavioral therapy groups, a cognitive behavioral stress intervention, a communication skills training package* combined with a couple-based intervention as well as a *behavioral therapy group* in comparison to the control group (no intervention course) on anxiety during pregnancy. *Individual structured psychotherapy* showed a significant improvement on state anxiety scores.

An *acceptance and commitment therapy (ACT)* intervention showed a positive effect on global distress, but must be interpreted carefully due to weak ratings. A

cognitive analytic therapy did not show any difference in trait/state anxiety between the randomized groups, but this result has to be interpreted carefully due to a weak rating.

Contrary results were shown regarding *biofeedback interventions*. While one study found that bio feedback significantly reduced FOC, another study did not find significant long-term improvements, but those results have to be interpreted carefully due to weak ratings.

Further, single studies about *calming virtual reality environments* (result only shown within qualitative data), a *motivational interviewing psychotherapy, emotional freedom techniques, breathing awareness, a mental health training course* (weak rating), the *group intervention Nyytti@* (with psychoeducation elements, the lifespan model of motivation, practices to support mentalisation and mind-body connection) (weak rating), *supportive care* through a midwife (physical, emotional, instructional, informational, advocacy support), *"lifestyle based education"*, different psychological interventions (like psychological support, education, relaxation training, family support, music listening) (weak rating), a *mind body intervention* (weak rating) and *group discussion* together with routine care seem to be effective in reducing anxiety during pregnancy and childbirth. While an *internet-based problem-solving treatment* did not show positive effects on reducing anxiety during pregnancy and antenatal class attendance reduced delivery fear significantly in first time, but not for mothers giving birth for the second or subsequent time.

Comparisons with other studies

Comparison of reviews based on mindfulness-based interventions: Compared to earlier systematic reviews within the literature, there are on the one side conflicting on the other side similar results. The systematic review and meta-analysis of Dhillon et al. also found benefits of mindfulness based interventions in RCT and non-RCT studies on anxiety, while mixed results are shown for perceived stress [27]. The pilot randomized trial of Beattie et al. in this present review did not find a beneficial effect of mindfulness based interventions on perceived stress [139].

The study of Hall et al. was not accessible for this review, as only the abstract exists, but the results seem to show a non-significant trend of mindfulness training towards a decrease in anxiety, this stands in conflict with the results within this present study [140-143].

Conflicting results also exist comparing the present review with the study of Lever and Taylor who in their between-group analysis did not find any significant effect of mindfulness based interventions on anxiety in comparison to control conditions. This is opposed to the results from RCTs (with moderate to strong ratings) of the present study, finding significant between group differences on those examined variables [29,103,112,125,126,129,130].

Consistent to the present review, Matvienko-Sikar et al. found in the majority of reviewed papers significant decreases of anxiety through mindfulness based interventions [30].

In the review of Riet et al. mixed results were found

in three studies regarding between-group effects of mindfulness based interventions on anxiety [31]. While two studies found a significant between-group effect in favor of the intervention group, one did not, but showed a significant decrease of anxiety due to mindfulness in the intervention group [31]. In the present study the between-group effect of mindfulness in moderate to strong RCTs on the outcome anxiety is significant and therefore shows a similar result compared to Riet et al. [31].

In the study of Shi and MacBeth seven RCTs showed significant reductions on anxiety due to mindfulness based interventions compared with control groups and four of five non-controlled studies also showed a significant decrease of anxiety [32]. The RCT results within this present review with anxiety as an outcome seem to match with this result.

Comparison of reviews based on hypnotherapeutic interventions: Two systematic reviews examined the effect of hypnosis based interventions and found a positive impact of hypnosis-based interventions on childbirth experience [21,22,42]. This finding fits the positive effect of hypnosis-based interventions on anxiety and fear found in the present study.

Comparison of reviews based on (psycho-) educational interventions: Akgün examined in their systematic review and meta-analysis the effect of psychoeducation on fear of childbirth and stated as a result, that fear of childbirth was reduced through psychoeducation [41]. This result also fits the result found within this present review, as positive effects of psychoeducation for fears are summarized. Only one study did not validate this result, but has to be interpreted carefully, due to weak ratings. The clinical review and meta-analysis of MoghaddamHosseini et al. found a significant effect of educational interventions on reducing fear of childbirth [21]. The systematic reviews of Striebich et al. supported this result and Stoll et al. also showed positive effects in reducing different anxiety/fear levels [23,25].

Comparison of reviews based on different therapeutic techniques/schools: Three reviews focussed on psychotherapy interventions [33-35]. Li et al. examined the effect of interpersonal psychotherapy (IPT) on fears and anxieties and found that interpersonal psychotherapy and peer supported interpersonal psychotherapy reduces fears and anxieties during pregnancy and childbirth [33]. Sockol et al. found similar effects of IPT on anxiety and fears in perinatal women [35]. Contrary to this outcome, the review of Ponting could not confirm this positive effect of IPT [34]. No study within this review focused on interpersonal psychotherapy.

Regarding cognitive behavioral therapy (studied was the general CBT, not mindfulness based), the systematic review of Li et al. found mixed effects [33]. While one study did not find a significant between-group effect, other studies found a significant reduction of fear and anxiety during pregnancy and childbirth [33,38]. Striebich et al. stated, that cognitive therapy sessions are effective in reducing fear of childbirth and van Ravesteyn found the same effect for anxiety

disorders [23,26]. Within this present study mixed results about internet based cognitive behavioral therapy are found, while Larsson et al. and Loughnan did not find a between-group effect for internet based cognitive behavioral therapy, Nieminen et al., Rondung et al. and Shahsavan et al. showed significant effects in favour of internet based cognitive behavioral therapy [38,83,87,89,91]. This result has to be interpreted carefully as the studies from Larsson et al., Loughnan and Nieminen et al. are rated as weak [83,87,127].

Based on the studies of the review of Li et al., Striebich et al. and the present study, mixed results for CBT are shown, but when taking the ratings of these studies into account, CBT seems to be effective in reducing fears and anxieties during pregnancy and childbirth [23,33].

Comparison of reviews based on writing therapy: Within this present systematic review, Montazeri et al. showed a significant and reductional effect of writing therapy sessions on anxiety during pregnancy, which opposes the result of the meta-analysis from Qian et al., which did not find a significant reducing effect of expressive writing therapy on anxiety [40,120].

Comparison with reviews based on e-health and technology-based interventions: The systematic review and meta-analysis of Bayrampour et al. demonstrated a significant reduction of anxiety scores in the e-health compared to a control group [36]. Another systematic review and meta-analysis showed mixed results for the effect of online cognitive behavioral therapy [38]. Mixed results about internet based cognitive behavioral therapy are also found within the present systematic review. For e-health interventions and their effect on reducing anxiety and fear regarding pregnancy and childbirth, mixed results are found within the present study and earlier systematic reviews and meta-analyses.

Limitations and Strengths

Limitations of the present systematic review are differences in the conceptualisations and operationalisations of prenatal anxiety and anxiety and fear during childbirth. Most of the studies are from Iran, so the results could be biased due to cultural background. Also the present review only included studies written in English and German. Another limitation is, that only one person rated the EPHPP criteria. The settings and duration of interventions differed between the studies and this fact could bias the results.

One strength is, that the present study until now includes the largest number of studies in systematic reviews addressing similar topics.

Study Implications

The present review only focused on certain therapeutic schools and psychotherapeutic interventions. Cognitive behavioral therapies are overrepresented, while there is only one psychoanalytic study and no study on systemic therapy, and its effect on fears during pregnancy and childbirth. Furthermore, there is a need for manualized therapeutic interventions, with regards to a combination of effective intervention components.

Conclusion

Within this systematic review, a wide range of psychological interventions are shown to be effective in reducing fears and anxieties during pregnancy and childbirth. These results are partly consistent with earlier systematic reviews and meta-analyses. Further research should address other acknowledged psychotherapeutic practices, like psychoanalytic or psychodynamic as well as systemic interventions, as they are underrepresented within this review. Furthermore, there is a need for manualized therapeutic interventions, with regards to a combination of effective intervention components.

Declarations

Ethics approval and consent to participate: Not applicable.

Consent for publication: I declare that all co-authors have given informed consent to publication of the manuscript.

Funding: We acknowledge that funding was kindly provided by the European Regional Development Fund, grant number: 85037910.

Competing interests: None declared.

Availability of data and materials: The Zotero library and the excel sheet to extract duplicates used in the search strategy are available from the corresponding author on reasonable request.

Authors' contribution: CB conceived the study and screened and selected the studies. GG independently checked the content. All authors helped draft the manuscript. CB and GG read and approved the final manuscript. CB is the lead author.

Acknowledgement: We are sincerely thankful to Julia Huwer, to support us during the course of this paper.

References

1. Madhavanprabhakaran GK, D'Souza MS, Nairy KS (2015) Prevalence of pregnancy anxiety and associated factors. *International Journal of Africa Nursing Sciences* 3: 1-7.
2. O'Connell Maeve A, Leahy-Warren P, Kenny LC, O'Neill SM, Khashan AS (2019) The prevalence and risk factors of fear of childbirth among pregnant women: A cross-sectional study in Ireland. *Acta Obstetrica et Gynecologica Scandinavica* 98: 1014-1023.
3. Adams S, Eberhard-Gran M, Eskild A (2012) Fear of childbirth and duration of labour: A study of 2206 women with intended vaginal delivery: Fear of childbirth and duration of labour. *BJOG: An International Journal of Obstetrics & Gynaecology* 119: 1238-1246.
4. Junge C, von Soest T, Weidner K, Seidler A, Eberhard-Gran M, et al. (2018) Labor pain in women with and without severe fear of childbirth: A population-based, longitudinal study. *Birth* 45: 469-477.
5. Laursen M, Johansen C, Hedegaard M (2009) Fear of childbirth and risk for birth complications in nulliparous women in the Danish National Birth Cohort. *BJOG: An International Journal of Obstetrics & Gynaecology* 116: 1350-1355.
6. Nettelblatt P, Fagerström CF, Uddenberg N (1976) The significance of reported childbirth pain. *Journal of Psychosomatic Research* 20: 215-221.
7. Reck C, Zimmer K, Dubber S, Zipser B, Schlehe B, et al. (2013) The influence of general anxiety and childbirth-specific anxiety on birth outcome. *Archives of Women's Mental Health* 16: 363-369.

8. Striebich S, Ayerle GM (2020) Große Angst vor der Geburt bei schwangeren Frauen in Deutschland: Erfahrungen und Orientierungen in Bezug auf die Schwangerenbetreuung und die Geburtsvorbereitung – eine rekonstruktive Studie. *GMS Zeitschrift für Hebammenwissenschaft*, 7.
9. Elvander C, Cnattingius S, Kjerulff KH (2013) Birth Experience in Women with Low, Intermediate or High Levels of Fear: Findings from the First Baby Study. *Birth* 40: 289-296.
10. Slade P, Balling K, Sheen K, Houghton G (2019) Establishing a valid construct of fear of childbirth: Findings from in-depth interviews with women and midwives. *BMC Pregnancy and Childbirth* 19: 96.
11. Haines HM, Rubertsson C, Pallant JF, Hildingsson I (2012) The influence of Women's fear, attitudes and beliefs of childbirth on mode and experience of birth. *BMC Pregnancy and Childbirth* 12: 55.
12. Räisänen S, Lehto S, Nielsen H, Gissler M, Kramer M, et al. (2014) Fear of childbirth in nulliparous and multiparous women: A population-based analysis of all singleton births in Finland in 1997-2010. *BJOG: An International Journal of Obstetrics & Gynaecology* 121: 965-970.
13. Söderquist J, Wijma B, Thorbert G, Wijma K (2009) Risk factors in pregnancy for post-traumatic stress and depression after childbirth. *BJOG: An International Journal of Obstetrics & Gynaecology* 116: 672-680.
14. Striebich S, Ayerle GM (2019) Fear of childbirth (FOC): Pregnant women's perceptions towards the impending hospital birth and coping resources – a reconstructive study. *Journal of Psychosomatic Obstetrics & Gynecology* 41: 231-239.
15. Leigh B, Milgrom J (2008) Risk factors for antenatal depression, postnatal depression and parenting stress. *BMC Psychiatry* 8: 24.
16. O'Connor TG, Heron J, Golding J, Beveridge M, Glover V (2002) Maternal antenatal anxiety and children's behavioural/emotional problems at 4 years. *British Journal of Psychiatry* 180: 502-508.
17. Sutter-Dallay AL, Giaccone-Marcasche V, Glatigny-Dallay E, Verdoux H (2004) Women with anxiety disorders during pregnancy are at increased risk of intense post-natal depressive symptoms: A prospective survey of the MATQUID cohort. *European Psychiatry* 19: 459-463.
18. Viswasam K, Eslick GD, Starcevic V (2019) Prevalence, onset and course of anxiety disorders during pregnancy: A systematic review and meta analysis. *Journal of Affective Disorders* 255: 27-40.
19. Mortazavi F, Mehrabadi M, KiaeeTabar R (2021) Pregnant women's well-being and worry during the COVID-19 pandemic: A cross-sectional study. *BMC Pregnancy and Childbirth* 21: 59.
20. Molgora S, Accordini M (2020) Motherhood in the Time of Coronavirus: The Impact of the Pandemic Emergency on Expectant and Postpartum Women's Psychological Well-Being. *Frontiers in Psychology* 11: 567155.
21. MoghaddamHosseini V, Nazarzadeh M, Jahanfar S (2018) Interventions for reducing fear of childbirth: A systematic review and meta-analysis of clinical trials. *Women and Birth* 31: 254-262.
22. Smith CA, Shewamene Z, Galbally M, Schmied V, Dahlen H (2019a) The effect of complementary medicines and therapies on maternal anxiety and depression in pregnancy: A systematic review and meta-analysis. *Journal of Affective Disorders* 245: 428-439.
23. Striebich S, Mattern E, Ayerle GM (2018) Support for pregnant women identified with fear of childbirth (FOC)/tokophobia – A systematic review of approaches and interventions. *Midwifery* 61: 97-115.
24. San Lazaro Campillo I, Meaney S, McNamara K, O'Donoghue K (2017) Psychological and support interventions to reduce levels of stress, anxiety or depression on women's subsequent pregnancy with a history of miscarriage: An empty systematic review. *BMJ Open* 7.
25. Stoll K, Swift EM, Fairbrother N, Nethery E, Janssen P (2018) A systematic review of nonpharmacological prenatal interventions for pregnancy-specific anxiety and fear of childbirth. *Birth* 45: 7-18.
26. van Ravesteyn LM, Lambregtse - van den Berg MP, Hoogendijk WJG, Kamperman AM (2017) Interventions to treat mental disorders during pregnancy: A systematic review and multiple treatment meta-analysis. *PLOS ONE* 12: e0173397.
27. Dhillon A, Sparkes E, Duarte RV (2017) Mindfulness-Based Interventions During Pregnancy: A Systematic Review and Meta-Analysis. *Mindfulness* 8: 1421-1437.
28. Hall H, Beattie J, Lau R, East C, Biro MA (2015) The effectiveness of mindfulness training on perinatal mental health: A systematic review. *Integrative Medicine Research* 4: 125.
29. Lever Taylor B, Cavanagh K, Strauss C (2016) The Effectiveness of Mindfulness-Based Interventions in the Perinatal Period: A Systematic Review and Meta-Analysis. *PLOS ONE* 11.
30. Matvienko-Sikar K, Lee L, Murphy G, Murphy L (2016) The effects of mindfulness interventions on prenatal well-being: A systematic review. *Psychology & Health* 31: 1415-1434.
31. Riet P, Francis L, Rees A (2020) Exploring the impacts of mindfulness and yoga upon childbirth outcomes and maternal health: An integrative review. *Scandinavian Journal of Caring Sciences* 34: 552-565.
32. Shi Z, MacBeth A (2017) The Effectiveness of Mindfulness-Based Interventions on Maternal Perinatal Mental Health Outcomes: A Systematic Review. *Mindfulness* 8: 823-847.
33. Li C, Sun X, Li Q, Sun Q, Wu B, Duan D (2020) Role of psychotherapy on antenatal depression, anxiety, and maternal quality of life. *Medicine* 99.
34. Ponting C, Mahrer NE, Zelcer H, Dunkel Schetter C, Chavira DA (2020) Psychological interventions for depression and anxiety in pregnant Latina and Black women in the United States: A systematic review. *Clinical Psychology & Psychotherapy* 27: 249-265.
35. Sockol LE (2018) A systematic review and meta-analysis of interpersonal psychotherapy for perinatal women. *Journal of Affective Disorders* 232: 316-328.
36. Bayrampour H, Trieu J, Tharmaratnam T (2019) Effectiveness of eHealth Interventions to Reduce Perinatal Anxiety: A Systematic Review and Meta-Analysis. *The Journal of Clinical Psychiatry* 80: 18r12386.
37. Lee Y, Cho S (2019) Technology-Supported Interventions for Pregnant Women: A Systematic Review. *CIN: Computers, Informatics, Nursing* 37: 501-512.
38. Loughnan SA, Joubert AE, Grierson A, Andrews G, Newby JM (2019) Internet-delivered psychological interventions for clinical anxiety and depression in perinatal women: A systematic review and meta-analysis. *Archives of Women's Mental Health* 22: 737-750.
39. Stratton E, Lampit A, Choi I, Calvo RA, Harvey SB, et al. (2017) Effectiveness of eHealth interventions for reducing mental health conditions in employees: A systematic review and meta-analysis. *PLOS ONE* 12: e0189904.
40. Qian J, Zhou X, Sun X, Wu M, Sun S, et al. (2020) Effects of expressive writing intervention for women's PTSD, depression, anxiety and stress related to pregnancy: A meta-analysis of randomized controlled trials. *Psychiatry Research* 288: 112933.
41. Akgün M, Boz İ, Özer Z (2019) The effect of psychoeducation on fear of childbirth and birth type: Systematic review and meta-analysis. *Journal of Psychosomatic Obstetrics & Gynecology* 41: 253-265.
42. Catsaros S, Wendland J (2020) Hypnosis-based interventions during pregnancy and childbirth and their impact on women's childbirth experience: A systematic review. *Midwifery* 84: 102666.
43. Bright KS, Charrois EM, Kashif Mughal M, Wajid A, McNeil D, et al. (2019) Interpersonal psychotherapy for perinatal women: A systematic review and meta-analysis protocol. *Systematic Reviews* 8: 9.
44. Brixval CS, Axelsen SF, Andersen SK, Due P, Koushede V (2014) The effect of antenatal education in small classes on obstetric and psychosocial outcomes: A systematic review and meta-analysis protocol. *Systematic Reviews* 3: 12.
45. O'Connell Maeve Anne, O'Neill SM, Dempsey E, Khashan AS, Leahy-Warren P, et al. (2019) Interventions for fear of childbirth (tocophobia). *Cochrane Database of Systematic Reviews* 5.
46. Smith CA, Shewamene Z, Galbally M, Schmied V, Dahlen H (2019) The effect of complementary medicines and therapies on maternal anxiety and depression in pregnancy: A systematic review and meta-analysis.

- Journal of Affective Disorders 245: 428-439.
47. Swift EM, Zoega H, Stoll K, Avery M, Gottfredsdóttir H (in press) Enhanced Ante-natal Care: Combining one-to-one and group Antenatal Care models to increase childbirth education and address childbirth fear. *Women and Birth* 34: 381-388.
 48. Abdollahi S, Faramarzi M, Delavar MA, Chehrazhi M, Gholinia H (2020) Effect of Psychotherapy on Reduction of Fear of Childbirth and Pregnancy Stress: A Randomized Controlled Trial. *Frontiers in Psychology* 11: 787.
 49. Areskog B (1982) Fear of Childbirth in Pregnant Women. *Advances in Psychosomatic Obstetrics and Gynecology*. Springer, Berlin Heidelberg.
 50. Zar M, Wijma K, Wijma B (2002) Relations between anxiety disorders and fear of childbirth during late pregnancy. *Clinical Psychology & Psychotherapy* 9: 122-130.
 51. Klabbbers GA (2016) Severe Fear of Childbirth: Its Features, Assessment, Prevalence, Determinants Consequences and Possible Treatments. *Psychological Topics* 25: 107-127.
 52. Størksen HT, Garthus-Niegel S, Adams SS, Vangen S, Eberhard-Gran M (2015) Fear of childbirth and elective caesarean section: A population-based study. *BMC Pregnancy and Childbirth* 15: 221.
 53. Wijma K (2003) Why focus on 'fear of childbirth'? *Journal of Psychosomatic Obstetrics & Gynecology* 24: 141-143.
 54. Poggi L, Goutaudier N, Séjourné N, Chabrol H (2018) When Fear of Childbirth is Pathological: The Fear Continuum. *Maternal and Child Health Journal* 22: 772-778.
 55. Hofberg K, Ward MR (2003) Fear of pregnancy and childbirth. *Postgraduate Medical Journal* 79: 505-510.
 56. Hofberg Kristina, Brockington I (2000) Tokophobia: An unreasoning dread of childbirth: A series of 26 cases. *British Journal of Psychiatry* 176: 83-85.
 57. Hofberg Kristina, Ward MR (2004) Fear of Childbirth, Tocophobia, and Mental Health in Mothers: The Obstetric-Psychiatric Interface: *Clinical Obstetrics and Gynecology* 47: 527-534.
 58. Bhatia MS, Jhanjee A (2012) Tokophobia: A dread of pregnancy. *Industrial Psychiatry Journal* 21: 158-159.
 59. O'Connell Maeve A, Leahy-Warren P, Khashan AS, Kenny LC, O'Neill SM (2017) Worldwide prevalence of tocophobia in pregnant women: Systematic review and meta-analysis. *Acta Obstetrica et Gynecologica Scandinavica* 96: 907-920.
 60. Wijma Klaas, Wijma B (2017) *A Woman Afraid to Deliver: How to Manage Childbirth Anxiety*. Bio-Psycho-Social Obstetrics and Gynecology. Springer International Publishing.
 61. Harrison V, Moore D, Lazard L (2020) Supporting perinatal anxiety in the digital age; a qualitative exploration of stressors and support strategies. *BMC Pregnancy and Childbirth* 20: 363.
 62. Misri S, Abizadeh J, Sanders S, Swift E (2015) Perinatal Generalized Anxiety Disorder: Assessment and Treatment. *Journal of Women's Health* 24: 762-770.
 63. Nasiri S, Akbari H, Tagharrobi L, Tabatabaee A (2018) The effect of progressive muscle relaxation and guided imagery on stress, anxiety, and depression of pregnant women referred to health centers. *Journal of Education and Health Promotion* 7.
 64. Kao MH, Hsu PF, Tien SF, Chen CP (2017) Effects of Support Interventions in Women Hospitalized With Preterm Labor. *Clinical Nursing Research* 28: 726-743.
 65. Jahdi F, Koohsarian Z, Rasouljan M, Montazerim A (2016) Evaluating the effectiveness of empowerment-based education on empowerment and anxiety in nulliparous women. *Acta Medica Mediterranea* 32: 1281-1287.
 66. Anton R, David D (2015) A randomized clinical trial of a new preventive rational emotive and behavioral therapeutical program of prepartum and postpartum emotional distress. *Journal of Evidence-Based Psychotherapies* 15: 3-15.
 67. Soltani F, Eskandari Z, Khodakarami B, Parsa P, Roshanaei G (2017) The effect of self-efficacy oriented counselling on controlling the fear of natural delivery in primi-gravida women. *Journal of Pharmaceutical Sciences and Research* 9: 1757-1761.
 68. Najafi Z, Mohebi S, Gharlipour Z, Rajabi M, Sharifirad G (2019) The effectiveness of emotional intelligence training on anxiety in primigravid women. *Cochrane Central Register of Controlled Trials* 8.
 69. Hennelly SE, Perman-Howe P, Foxcroft DR, Smith LA (2020) The feasibility of 'Mind the Bump': A mindfulness based maternal behaviour change intervention. *Complementary Therapies in Clinical Practice* 40: 101178.
 70. Airo (Toivanen) R, Korja R, Saisto T, Rouhe H, Muotka J, et al. (2018) Changes in emotions and personal goals in primiparous pregnant women during group intervention for fear of childbirth. *Journal of Reproductive and Infant Psychology* 36: 363-380.
 71. Amiri P, Mirghafourvand M, Esmailpour K, Kamalifard M, Ivanbagha R (2019) The effect of distraction techniques on pain and stress during labor: A randomized controlled clinical trial. *BMC Pregnancy and Childbirth* 19: 534.
 72. Andaroon N, Kordi M, Kimiaei S, Esmaeily H (2017) The effect of individual counseling program by a midwife on fear of childbirth in primiparous women. *Journal of Education and Health Promotion* 6: 97.
 73. Boryri T, Navidian A, Marghzari N (2018) Comparison of the Effect of Muscle Relaxation and Guided Imagery on Happiness and Fear of Childbirth in Primiparous Women Admitted to Health Care Centers. *International Journal of Women's Health and Reproduction Sciences* 7: 490-495.
 74. Boz İ, Akgün M, Duman F (2020) A feasibility study of a psychoeducation intervention based on Human Caring Theory in nulliparous women with fear of childbirth. *Journal of Psychosomatic Obstetrics & Gynecology*.
 75. Firouzan L, Kharaghani R, Zenoozian S, Moloodi R, Jafari E (2020) The effect of midwifery led counseling based on Gamble's approach on childbirth fear and self efficacy in nulligravida women. *BMC Pregnancy and Childbirth*, 20: 522.
 76. Haapio S, Kaunonen M, Arffman M, Åstedt-Kurki P (2017) Effects of extended childbirth education by midwives on the childbirth fear of first-time mothers: An RCT. *Scandinavian Journal of Caring Sciences* 31: 293-301.
 77. Hildingsson I, Rubertsson C, Karlström A, Haines H (2019) A known midwife can make a difference for women with fear of childbirth: birth outcome and women's experiences of intrapartum care. *Sexual & Reproductive Healthcare* 21: 33-38.
 78. Irmak Vural P, Aslan E (2019) Emotional freedom techniques and breathing awareness to reduce childbirth fear: A randomized controlled study. *Complementary Therapies in Clinical Practice* 35: 224-231.
 79. Kacperczyk-Bartnik J, Bartnik P, Symonides A, Sroka-Ostrowska N, Dobrowolska-Redo A, et al. (2019) Association between antenatal classes attendance and perceived fear and pain during labour. *Taiwanese Journal of Obstetrics and Gynecology* 58: 492-496.
 80. Karabulut Ö, Coşkuner Potur D, Doğan Merih Y, Cebeci Mutlu S, Demirci N (2016) Does antenatal education reduce fear of childbirth?: Does antenatal education reduce fear of childbirth. *International Nursing Review* 63: 60-67.
 81. Klabbbers GA, Wijma K, Paarlberg KM, Emons WHM, Vingerhoets AJJM (2019) Haptotherapy as a new intervention for treating fear of childbirth: A randomized controlled trial. *Journal of Psychosomatic Obstetrics & Gynecology* 40: 38-47.
 82. Kordi M, Bakhshi mohaddeseh, Masoudi S, Esmaily H (2017) The effect of childbirth psychoeducational program on severity childbirth fear in primigravida women. *Evidence Based Care* 7: 26-34.
 83. Larsson B, Karlström A, Rubertsson C, Ternström E, Ekdahl J, et al. (in press) Birth preference in women undergoing treatment for childbirth fear: A randomised controlled trial. *Women and Birth* 30: 460-467.
 84. Mary AJK, Latheef F, Vijayaraghavan R (2017) Effectiveness of selected mind body interventions on anxiety related to childbirth and labour

- outcomes. *Indian Journal of Traditional Knowledge* 16: 122-128.
85. Munkhondya BMJ, Munkhondya TE, Chirwa E, Wang H (2020) Efficacy of companion integrated childbirth preparation for childbirth fear, self-efficacy, and maternal support in primigravid women in Malawi. *BMC Pregnancy and Childbirth* 20: 48.
 86. van Y, Shinohara H, Kodama H (2018) Resting Heart Rate Variability and the Effects of Biofeedback Intervention in Women with Low-Risk Pregnancy and Prenatal Childbirth Fear. *Applied Psychophysiology and Biofeedback* 43: 113-121.
 87. Nieminen K, Andersson G, Wijma B, Ryding EL, Wijma K (2016) Treatment of nulliparous women with severe fear of childbirth via the Internet: A feasibility study. *Journal of Psychosomatic Obstetrics & Gynecology* 37: 37-43.
 88. Rahmani R, Yaghoobi Moghadam F, Hadizadeh-Talasz F, Rahmani MR (2019) The Effect of Peer Education on Fear of Childbirth in Pregnant Women: A Randomized Clinical Trial. *International Journal of Women's Health and Reproduction Sciences* 8: 209-214.
 89. Rondung E, Ternström E, Hildingsson I, Sundin Ö, Ekdahl J, et al. (2018) Comparing Internet-Based Cognitive Behavioral Therapy With Standard Care for Women With Fear of Birth: Randomized Controlled Trial. *JMIR Mental Health* 5: e10420.
 90. Rouhe H, Salmela-Aro K, Toivanen R, Tokola M, Halmesmäki E, et al. (2015) Group psychoeducation with relaxation for severe fear of childbirth improves maternal adjustment and childbirth experience – a randomised controlled trial. *Journal of Psychosomatic Obstetrics & Gynecology* 36: 1-9.
 91. Shahsavan F, Akbari N, Gharraee B, Abolghasemi J, Khedmat L (2020) The effect of internet-based guided self-help cognitive-behavioral therapies on Iranian women's psychological symptoms and preferred method of childbirth. *Perspectives in Psychiatric Care* 57: 138-147.
 92. Uçar T, Golbasi Z (2019) Effect of an educational program based on cognitive behavioral techniques on fear of childbirth and the birth process. *Journal of Psychosomatic Obstetrics & Gynecology* 40: 146-155.
 93. Abbasi P, Mohammad-Alizadeh Charandabi S, Mirghafourvand M (2020) Comparison of the effect of educational software and booklet on anxiety and pain during labour: A randomised controlled clinical trial. *Journal of Obstetrics and Gynaecology* 41: 1-8.
 94. Ekrami F, Mohammad-Alizadeh Charandabi S, Babapour Kheiroddin J, Mirghafour-vand M (2019) The Effect of Counselling on Depression and Anxiety of Women with Unplanned Pregnancy: A Randomized Controlled Trial. *Community Mental Health Journal* 55: 1047-1056.
 95. Fontein-Kuipers YJ, Ausems M, de Vries R, Nieuwenhuijze MJ (2016) The effect of Wazzup Mama?! An antenatal intervention to prevent or reduce maternal distress in pregnancy. *Archives of Women's Mental Health* 19: 779-788.
 96. Goetz M, Schiele C, Müller M, Matthies LM, Deutsch TM, et al. (2020) Effects of a Brief Electronic Mindfulness-Based Intervention on Relieving Prenatal Depression and Anxiety in Hospitalized High-Risk Pregnant Women: Exploratory Pilot Study. *Journal of Medical Internet Research* 22: 16.
 97. Hamilton J, Saxon D, Best E, Glover V, Walters SJ, et al. (2020) A randomized, controlled pilot study of cognitive analytic therapy for stressed pregnant women with underlying anxiety and depression in a routine health service setting. *Clinical Psychology & Psychotherapy* 28: 394-408.
 98. Parsa P, Saeezadeh N, Zahra Masoumi S, Roshanaei G (2016) The Effectiveness of Counseling in Reducing Anxiety Among Nulliparous Pregnant Women. *Journal of Family and Reproductive Health* 10: 198-205.
 99. Rajeswari S, Sanjeeva Reddy N (2020) Efficacy of progressive muscle relaxation on pregnancy outcome among anxious Indian primi mothers. *Iranian Journal of Nursing and Midwifery Research* 25: 23.
 100. Salehi F, Pourasghar M, Khalilian A, Shahhosseini Z (2016) Comparison of group cognitive behavioral therapy and interactive lectures in reducing anxiety during pregnancy: A quasi experimental trial. *Medicine* 95: e5224.
 101. Sanaati F, Mohammad-Alizadeh Charandabi S, Farrokh Eslamlo H, Mirghafourvand M, Alizadeh Sharajabad, F (2017) The effect of lifestyle-based education to women and their husbands on the anxiety and depression during pregnancy: A randomized controlled trial. *The Journal of Maternal-Fetal & Neonatal Medicine* 30: 870-876.
 102. Toosi M, Akbarzadeh M, Ghaemi Z (2017) The Effect of Relaxation on Mother's Anx-iety and Maternal-Fetal Attachment in Primiparous IVF Mothers. *Journal of the National Medical Association* 109: 164-171.
 103. Zhang JY, Cui YX, Zhou YQ, Li YL (2019) Effects of mindfulness-based stress reduction on prenatal stress, anxiety and depression. *Psychology, Health & Medicine* 24: 51-58.
 104. Esfandiari M, Faramarzi M, Nasiri-Amiri F, Parsian H, Chehrizi M, et al. (2020) Effect of supportive counseling on pregnancy-specific stress, general stress, and prenatal health behaviors: A multicenter randomized controlled trial. *Patient Education and Counseling* 103: 2297-2304.
 105. Legrand F, Grévin-Laroche C, Josse E, Polidori G, Quinart H, et al. (2017) Effects of hypnosis during pregnancy: A psychophysiological study on maternal stress. *Medical Hypotheses* 102: 123-127.
 106. Mirtabar SM, Faramarzi M, Khazaei R, Dini M (2020) Efficacy of psychotherapy for anxiety reduction in hospital management of women successfully treated for preterm labor: A randomized controlled trial. *Women & Health* 60: 1151-1163.
 107. Sheikh-Azadi E, Bakhteh A, Rezaei M, Ashtareian H (2016) The effect of group discussion in primigravidas. *Research Journal of Medical Sciences* 10: 498-502.
 108. Aksoy Derya Y, Altiparmak S, Akça E, Gökbulut N, Yilmaz AN (2021) Pregnancy and birth planning during COVID-19: The effects of tele-education offered to pregnant women on prenatal distress and pregnancy-related anxiety. *Midwifery* 92: 102877.
 109. Bazrafshan S, Kheirkhah M, Inanlou M, Rasouli M (2020) Controlling the anxiety in Iranian pregnant women at risk of preterm labor by undergoing the counseling group intervention. *Journal of Family Medicine and Primary Care* 9: 4016.
 110. Karamoozian M, Askarizadeh G (2015) Impact of Prenatal Cognitive-Behavioral Stress Management Intervention on Maternal Anxiety and Depression and Newborns' Apgar Scores. *Iranian Journal of Neonatology* 6: 14-23.
 111. Khojasteh F, Rezaee N, Safarzadeh A, Sahlabadi R, Shahrakipoor M (2016) Comparison of the Effects of Massage Therapy and Guided Imagery on Anxiety of Nulliparous Women during Pregnancy. *Der Pharmacia Lettre* 8: 1-7.
 112. Zarenejad M, Yazdkhasti M, Rahimzadeh M, Mehdizadeh Tourzani Z, Esmaelzadeh-Saeieh S (2020) The effect of mindfulness-based stress reduction on maternal anxiety and self-efficacy: A randomized controlled trial. *Brain and Behavior* 10: e01561
 113. Akbarian Z, Kohan S, Nasiri H, Ehsanpour S (2018) The Effects of Mental Health Training Program on Stress, Anxiety, and Depression during Pregnancy. *Iranian Journal of Nursing and Midwifery Research* 23: 93-97.
 114. Alipour Z, Kazemi A, Kheirabadi G, Eslami AA (2020) Marital communication skills training to promote marital satisfaction and psychological health during pregnancy: A couple focused approach. *Reproductive Health* 17: 23.
 115. Aslami E, Alipour A, Najib FS, Aghayosefi A (2016) A Comparative Study of Mindfulness Efficiency Based on Islamic-Spiritual Schemes and Group Cognitive Behavioral Therapy on Reduction of Anxiety and Depression in Pregnant Women 5: 144-152.
 116. Beevi Z, Low WY, Hassan J (2016) Impact of Hypnosis Intervention in Alleviating Psychological and Physical Symptoms During Pregnancy. *American Journal of Clinical Hypnosis* 58: 368-382.
 117. Hajmohamadi N, Ghalichi F, Bakhtari Aghdam F, Matlabi H (2018) The "Cooperative Supportive" Intervention for Improving Mental Health Status among Pregnant Women. *Journal of Caring Sciences* 7: 101-106.

118. Heller HM, Hoogendoorn AW, Honig A, Broekman BFP, van Straten A (2020) The Effectiveness of a Guided Internet-Based Tool for the Treatment of Depression and Anxiety in Pregnancy (MamaKits Online): Randomized Controlled Trial. *Journal of Medical Internet Research* 22: e15172.
119. Mohammadi MM, Parandin S (2019) Effect of the combination of Benson's relaxation technique and brief psychoeducational intervention on multidimensional pain and negative psychological symptoms of pregnant women: A randomized controlled trial. *Journal of Education and Health Promotion* 8: 91.
120. Montazeri M, Esmaeilpour K, Mohammad-Alizadeh-Charandabi S, Golizadeh S, Mirghafourvand M (2020) The Effect of Writing Therapy on Anxiety in Pregnant Women: A Randomized Controlled Trial. *Iranian Journal of Psychiatry and Behavioral Sciences* 14: e98256.
121. Sobhani E, Babakhani N, Alboe MR (2019) The Effectiveness of Mindfulness-Based Stress Reduction on the Depression, Anxiety, Stress, and Pain Perception in Females with Obstructed Labour-Induced Chronic Low Back Pain. *Iranian Journal of Psychiatry and Clinical Psychology* 25: 266-277.
122. Sridhar A, Shiliang Z, Woodson R, Kwan L (2020) Non-pharmacological anxiety reduction with immersive virtual reality for first-trimester dilation and curettage: A pilot study. *The European Journal of Contraception & Reproductive Health Care* 25: 480-483.
123. van der Zwan J, Huizink A, Lehrer P, Koot H, de Vente W (2019) The Effect of Heart Rate Variability Biofeedback Training on Mental Health of Pregnant and Non-Pregnant Women: A Randomized Controlled Trial. *International Journal of Environmental Research and Public Health* 16: 1051.
124. Yazdanimehr R, Omid A, Sadat Z, Akbari H (2016) The Effect of Mindfulness-integrated Cognitive Behavior Therapy on Depression and Anxiety among Pregnant Women: A Randomized Clinical Trial. *Journal of Caring Sciences* 5: 195-204.
125. Zhang M (2019) Effectiveness of mindfulness-based cognitive therapy for comorbid depression and anxiety in pregnancy: A randomized controlled trial. *Arch Womens Ment Health* 23: 51-59.
126. Krusche A, Dymond M, Murphy SE, Crane C (2018) Mindfulness for pregnancy: A randomised controlled study of online mindfulness during pregnancy. *Midwifery* 65: 51-57.
127. Loughnan SA, Sie A, Hobbs MJ, Joubert AE, Smith J, et al. (2019) A randomized controlled trial of "MUMentum Pregnancy": Internet-delivered cognitive behavioral therapy program for antenatal anxiety and depression. *Journal of Affective Disorders* 243: 381-390.
128. Price C, Kantrowitz-Gordon I, Calhoun R (2019) A pilot feasibility study of mindfulness childbirth education for women with a history of sexual trauma. *Complementary Therapies in Clinical Practice* 37: 102-108.
129. Warriner S, Crane C, Dymond M, Krusche A (2018) An evaluation of mindfulness-based childbirth and parenting courses for pregnant women and prospective fathers/partners within the UK NHS (MBCP-4-NHS). *Midwifery* 64: 1-10s.
130. Yang M, Jia G, Sun S, Ye C, Zhang R, et al. (2019) Effects of an Online Mindfulness Intervention Focusing on Attention Monitoring and Acceptance in Pregnant Women: A Randomized Controlled Trial. *Journal of Midwifery & Women's Health* 64: 68-77.
131. Noorbala AA, Afzali HM, Abedinia N, Akhbari M, Moravveji SA, et al. (2019) Investigation of the effectiveness of psychiatric interventions on the mental health of pregnant women in Kashan City - Iran: A clinical trial study. *Asian Journal of Psychiatry* 46: 79-86.
132. Waters CS, Annear B, Flockhart G, Jones I, Simmonds JR, et al. (2020) Acceptance and Commitment Therapy for perinatal mood and anxiety disorders: A feasibility and proof of concept study. *British Journal of Clinical Psychology* 59: 461-479.
133. Kang CM (2020) Effects of psychological intervention and relevant influence factors on pregnant women undergoing interventional prenatal diagnosis. *Journal of the Chinese Medical Association* 83: 202-205.
134. Uludağ E, Mete S (2020) The effect of nursing care provided based on the philosophy of hypnobirthing on fear, pain, duration, satisfaction and cost of labor: A single-blind randomized controlled study: The effect of nursing care on the labor process and cost. *Health Care for Women International* 12: 1-13.
135. Waisblat V, Langholz B, Bernard FJ, Arnould M, Benassi A, et al. (2017) Impact of a Hypnotically-Based Intervention on Pain and Fear in Women Undergoing Labor. *International Journal of Clinical and Experimental Hypnosis* 65: 64-85.
136. Duncan LG, Cohn MA, Chao MT, Cook JG, Riccobono J, et al. (2017) Benefits of preparing for childbirth with mindfulness training: A randomized controlled trial with active comparison. *BMC Pregnancy and Childbirth* 17: 140.
137. Seyed Kaboli K, Mahmoodi Z, Mehdizadeh Tourzani Z, Tehranizadeh M, Kabir K, et al. (2017) The Effect of Group Counseling Based on Cognitive-Behavioral Approach on Pregnancy-Specific Stress and Anxiety. *Shiraz E-Medical Journal* 18: e13183.
138. İsbir GG, Serçekuş P (2015) The Effects of Intrapartum Supportive Care on Fear of Delivery and Labor Outcomes: A Single-Blind Randomized Controlled Trial. *Journal of Nursing Research* 25: 112-119.
139. Beattie J, Hall H, Biro MA, East C, Lau R (2017) Effects of mindfulness on maternal stress, depressive symptoms and awareness of present moment experience: A pilot randomised trial. *Midwifery* 50: 174-183.
140. Hall WA, Stoll K, Hutton EK, Brown H (2012) A prospective study of effects of psychological factors and sleep on obstetric interventions, mode of birth, and neonatal outcomes among low-risk British Columbian women. *BMC Pregnancy and Childbirth* 12: 78.
141. Moher D, Liberati A, Tetzlaff J, Altman DG, The PRISMA Group (2009) Preferred Reporting Items for Systematic Reviews and Meta-Analyses: The PRISMA Statement. *PLoS Medicine* 6: e1000097.
142. Moudi Z, Molashahi B, Ansari H, Imani M (2018) The effect of a care program and social support on anxiety level in mothers of late preterm infants in Sistan and Baluchestan Iran. *Women & Health* 59: 569-578.
143. Zhang A, Park S, Sullivan JE, Jing S (2018) The Effectiveness of Problem-Solving Therapy for Primary Care Patients' Depressive and/or Anxiety Disorders: A Systematic Review and Meta-Analysis. *The Journal of the American Board of Family Medicine* 31: 139-150.
144. National Collaboration Center for Methods and Tools (2020) Quality Assessment Tool for Quantitative Studies.